

Semi-infinite Lakshmibai-Seshadri path model for level-zero extremal weight modules over quantum affine algebras^{*}

Motohiro Ishii

Research Center for Pure and Applied Mathematics,
Graduate School of Information Sciences, Tohoku University,
Aramaki aza Aoba 6-3-09, Aoba-ku, Sendai 980-8579, Japan
(e-mail: ishii@math.is.tohoku.ac.jp)

Satoshi Naito

Department of Mathematics, Tokyo Institute of Technology,
2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan
(e-mail: naito@math.titech.ac.jp)

Daisuke Sagaki

Institute of Mathematics, University of Tsukuba,
Tsukuba, Ibaraki 305-8571, Japan
(e-mail: sagaki@math.tsukuba.ac.jp)

Abstract

We introduce semi-infinite Lakshmibai-Seshadri paths by using the semi-infinite Bruhat order (or equivalently, Lusztig's generic Bruhat order) on affine Weyl groups in place of the usual Bruhat order. These paths enable us to give an explicit realization of the crystal basis of an extremal weight module of an arbitrary level-zero dominant integral extremal weight over a quantum affine algebra. This result can be thought of as a full generalization of our previous result (which uses Littelmann's Lakshmibai-Seshadri paths), in which the level-zero dominant integral weight is assumed to be a positive-integer multiple of a level-zero fundamental weight.

1 Introduction.

In our previous papers [NS1, NS3], we gave a combinatorial realization of the crystal basis $\mathcal{B}(m_i\varpi_i)$ of the extremal weight module $V(m_i\varpi_i)$ of extremal weight $m_i\varpi_i$, where $m_i \in \mathbb{Z}_{\geq 1}$

^{*}2010 Mathematics Subject Classification. Primary: 17B37, Secondary: 17B67, 81R50, 81R10.

and ϖ_i is the i -th level-zero fundamental weight for $i \in I$, over the quantum affine algebra $U_q(\mathfrak{g}_{\text{af}})$ in terms of Lakshmibai-Seshadri (LS for short) paths of shape $m_i \varpi_i$ in the sense of [Li2]; however, in [NS4], we showed that it is impossible to give a realization of the crystal basis $\mathcal{B}(\lambda)$ of the extremal weight module $V(\lambda)$ of a general level-zero dominant integral extremal weight λ in terms of Littelmann's LS paths of shape λ . The purpose of this paper is to overcome this difficulty, and to give an explicit realization of the crystal basis of the extremal weight module $V(\lambda)$ for a level-zero dominant integral weight λ in full generality; however, we assume that an affine Lie algebra $\mathfrak{g}_{\text{af}}$ is of untwisted type throughout this paper.

Extremal weight modules over the quantized universal enveloping algebras of symmetrizable Kac-Moody algebras were introduced by Kashiwara [Kas1] in his study of the level-zero part of the modified quantized universal enveloping algebra (see [Lu2]) of an affine Lie algebra.

Let λ be an integral weight for an affine Lie algebra $\mathfrak{g}_{\text{af}}$. If λ is of positive (resp., negative) level, then the extremal weight module $V(\lambda)$ is just the integrable highest (resp., lowest) weight module over $U_q(\mathfrak{g}_{\text{af}})$. However, in the case when λ is of level-zero, the structure of $V(\lambda)$ is much more complicated than in the case of positive or negative level. In fact, it is known ([N2, Remark 2.15]; see also [CP, Proposition 4.5]) that $V(\lambda)$ is isomorphic to the quantum Weyl module $W_q(\lambda)$ introduced by Chari and Pressley ([CP]).

Also, in the case when $\mathfrak{g}_{\text{af}}$ is an untwisted affine Lie algebra of type A , D , or E , an extremal weight module can be thought of as a universal standard module. Here standard modules M_P , parametrized by Drinfeld polynomials P , were constructed by Nakajima ([N1]) by use of quiver varieties, as a new basis of the Grothendieck ring $\text{Rep } U_q(L\mathfrak{g})$ of finite-dimensional modules (of type 1) over the quantum loop algebra $U_q(L\mathfrak{g})$, where $\mathfrak{g} \subset \mathfrak{g}_{\text{af}}$ is the canonical finite-dimensional subalgebra (of type A , D , or E), and $L\mathfrak{g} = \mathbb{C}[t, t^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}$; unique irreducible quotients L_P of the standard modules M_P form another basis of $\text{Rep } U_q(L\mathfrak{g})$.

More precisely, for a level-zero dominant integral weight $\lambda = \sum_{i \in I} m_i \varpi_i$ with $m_i \in \mathbb{Z}_{\geq 0}$, $i \in I$, the universal standard module $M(\lambda)$ is defined to be the Grothendieck group $K^{G_\lambda \times \mathbb{C}^*}(\mathfrak{L}(\lambda))$ of $G_\lambda \times \mathbb{C}^*$ -equivariant coherent sheaves on a certain Lagrangian subvariety $\mathfrak{L}(\lambda)$ of the quiver variety $\mathfrak{M}(\lambda)$, where $G_\lambda := \prod_{i \in I} GL_{m_i}(\mathbb{C})$. For an I -tuple $P = (P_i(u))_{i \in I}$ of monic polynomials with coefficients in $\mathbb{C}(q)$ such that $\deg P_i(u) = m_i$ (called a Drinfeld polynomial), the corresponding standard module M_P is obtained from $M(\lambda)$ as the specialization $M(\lambda) \otimes_{R(G_\lambda)[q, q^{-1}]} \mathbb{C}(q)$, where $R(G_\lambda) \cong \bigotimes_{i \in I} \mathbb{Z}[x_{i,1}^{\pm 1}, \dots, x_{i,m_i}^{\pm 1}]^{\mathfrak{S}_{m_i}}$ is the representation ring of G_λ , and by the algebra homomorphism $R(G_\lambda)[q, q^{-1}] \rightarrow \mathbb{C}(q)$, the indeterminates $x_{i,1}, \dots, x_{i,m_i}$ are sent to the roots of the polynomial $P_i(u)$ for $i \in I$. In [N2, Theorem 2], Nakajima proved that there exists a $U_q(\mathfrak{g}'_{\text{af}}) \otimes_{\mathbb{Z}[q, q^{-1}]} R(G_\lambda)[q, q^{-1}]$ -module isomorphism between the extremal weight module $V(\lambda)$ and $M(\lambda) \otimes_{\mathbb{Z}[q, q^{-1}]} \mathbb{C}(q)$, as a by-product of his proof of Kashiwara's conjecture (see [Kas5, §13]) on the structure of extremal weight modules of level-zero extremal weights, where $\mathfrak{g}'_{\text{af}} = (\mathbb{C}[t, t^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}) \oplus \mathbb{C}c$.

Now, let λ be a level-zero dominant integral weight of the form $\lambda = \sum_{i \in I} m_i \varpi_i$, with $m_i \in \mathbb{Z}_{\geq 0}$ for all $i \in I$. In [BN], Beck and Nakajima proved Kashiwara's conjecture above for all affine Lie algebras, and in particular, showed that there exists an isomorphism of crystals

$$\mathcal{B}(\lambda) \cong \bigotimes_{i \in I} \mathcal{B}(m_i \varpi_i). \quad (1)$$

Soon afterward, in [NS1, NS3], we gave a combinatorial realization of the crystal basis $\mathcal{B}(m_i \varpi_i)$ for each $i \in I$ in terms of LS paths of shape $m_i \varpi_i$. Namely, we proved that the set $\mathbb{B}(m_i \varpi_i)$ of LS paths of shape $m_i \varpi_i$ is isomorphic as a crystal to $\mathcal{B}(m_i \varpi_i)$, though neither of (the crystal graphs of) these two crystals is connected if $m_i > 1$. However, it turned out in [NS4] that the crystals $\mathbb{B}(\lambda)$ and $\mathcal{B}(\lambda)$ are not necessarily isomorphic for a general level-zero dominant integral weight λ ; for example, if λ is of the form $\sum_{i \in K} \varpi_i$ for $K \subset I$ with $\#K \geq 2$, then the crystals $\mathbb{B}(\lambda)$ and $\mathcal{B}(\lambda)$ are never isomorphic, though both of these crystals are connected (for details, see [NS4, Appendix]). This is mainly because each connected component of the crystal $\mathbb{B}(\lambda)$ has fewer extremal elements than that of the crystal $\mathcal{B}(\lambda)$ has.

Let us explain the situation above more precisely. As a consequence of the isomorphism (1), we see (Proposition 5.1.1) that the set $\{u_x := xu_\lambda \mid x \in W_{\text{af}}\}$ of extremal elements in the connected component $\mathcal{B}_0(\lambda)$ of $\mathcal{B}(\lambda)$ containing the extremal element u_λ of extremal weight λ is in bijective correspondence with the quotient set $W_{\text{af}}/(W_J)_{\text{af}}$, where $W_{\text{af}} = W \ltimes \{t_\xi \mid \xi \in Q^\vee\}$ is the (affine) Weyl group of $\mathfrak{g}_{\text{af}}$, and $(W_J)_{\text{af}} := W_J \ltimes \{t_\xi \mid \xi \in Q_J^\vee\}$, with $J := \{i \in I \mid m_i = 0\}$, is identical to the stabilizer $\{x \in W_{\text{af}} \mid xu_\lambda = u_\lambda\}$ of u_λ in W_{af} . In contrast, the set $\{\pi_{x\lambda} := (x\lambda; 0, 1) \mid x \in W_{\text{af}}\}$ of extremal elements in the connected component $\mathbb{B}_0(\lambda)$ of $\mathbb{B}(\lambda)$ containing the straight line path $\pi_\lambda := (\lambda; 0, 1)$ of weight λ is in bijective correspondence with the quotient set $W_{\text{af}}/(W_{\text{af}})_\lambda$, where $(W_{\text{af}})_\lambda$ is the stabilizer of λ in W_{af} , and is identical to $W_J \ltimes \{t_\xi \mid \langle \xi, \lambda \rangle = 0\}$. Here, we have $(W_J)_{\text{af}} \subset (W_{\text{af}})_\lambda$ in general, with equality if and only if λ is a nonnegative integer multiple of ϖ_i for some $i \in I$.

In order to overcome the difficulty mentioned above, we introduce the notion of semi-infinite Lakshmibai-Seshadri (SiLS for short) paths of shape λ . A SiLS path of shape λ is, by definition, a pair $(\mathbf{x}; \mathbf{a})$ of a decreasing sequence $\mathbf{x} : x_1 >_{\frac{\infty}{2}} x_2 >_{\frac{\infty}{2}} \cdots >_{\frac{\infty}{2}} x_s$ in the set $(W^J)_{\text{af}}$ of Peterson's coset representatives for the cosets in $W_{\text{af}}/(W_J)_{\text{af}}$, equipped with the semi-infinite Bruhat order $\geq_{\frac{\infty}{2}}$, and an increasing sequence $\mathbf{a} : 0 = a_0 < a_1 < \cdots < a_s = 1$ of rational numbers, while a (usual) LS path of shape λ is a pair $(\boldsymbol{\lambda}; \mathbf{a})$ of a decreasing sequence $\boldsymbol{\lambda} : \lambda_1 > \lambda_2 > \cdots > \lambda_s$ of elements in the affine Weyl group orbit $W_{\text{af}}\lambda$ through λ , equipped with the partial order $\geq$ which Littelmann defined in [Li2], and an increasing sequence $\mathbf{a}$ of rational numbers as above.

The coset representatives $(W^J)_{\text{af}}$ were originally introduced by Peterson ([P]; see also [LS]) in his study of the relationship between the T -equivariant homology (ring) of the affine Grassmannian $G(\mathbb{C}((t)))/G(\mathbb{C}[[t]])$ and the T -equivariant (small) quantum cohomology ring $QH_T^\bullet(G/P)$ of the partial flag variety G/P , where G denotes a simply-connected simple algebraic group over

$\mathbb{C}$, $P \subset G$ a parabolic subgroup, and $T \subset G$ a maximal torus. Also, we see from [S, Claim 4.14] that the semi-infinite Bruhat order on the affine Weyl group W_{af} is a slight modification of Lusztig's generic Bruhat order on W_{af} (for details, see Appendix A); the generic Bruhat order was originally introduced by Lusztig ([Lu1]) in his study of the conjectural character formula for the irreducible quotient of the Weyl module of a simply-connected almost simple algebraic group over an algebraically closed field of positive characteristic. As for the geometric meaning of the semi-infinite Bruhat order, it is known ([FFKM, §5]; see also [FF, §4]) that the semi-infinite Bruhat order describes the closure relation among the fine Schubert strata, parametrized by W_{af} , of the Drinfeld compactification of the variety of algebraic maps of a fixed degree from the complex projective line $\mathbb{P}^1$ to the flag variety G/B . In addition, we remark that in Peterson's lecture note ([P]), the semi-infinite Bruhat order (or, stable Bruhat order in his terminology) plays an important role, and that some of our arguments in the study of SiLS paths use (parabolic) quantum Bruhat graphs ([BFP, LNS³1]), which appear in the equivariant quantum Chevalley formula for $QH_T^\bullet(G/P)$ ([Mi]).

Now we are ready to state our main results. First, we prove that the natural surjection from the poset $(W^J)_{\text{af}}$ (equipped with the semi-infinite Bruhat order) onto the poset $W_{\text{af}}\lambda$ (equipped with Littelmann's partial order) is order-preserving, and hence that there exists a surjection from the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ of SiLS paths of shape λ onto the set $\mathcal{B}(\lambda)$ of LS paths of shape λ (Proposition 3.1.3). Next, we define a crystal structure on $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ for $U_q(\mathfrak{g}_{\text{af}})$ in such a way that this surjection becomes a morphism of crystals (Theorem 3.1.5). Then, each connected component of the resulting crystal $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ indeed has as many extremal elements as that of the crystal $\mathcal{B}(\lambda)$ has (Proposition 3.2.2). Also, we can prove that the crystal $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ has as many connected components as the crystal $\mathcal{B}(\lambda)$ has (Proposition 3.2.4). Combining the results above, we finally obtain the following (Theorem 3.2.1).

Theorem. *Let $\mathfrak{g}_{\text{af}}$ be an untwisted affine Lie algebra, and $\lambda = \sum_{i \in I} m_i \varpi_i$ a level-zero dominant integral weight, with $m_i \in \mathbb{Z}_{\geq 0}$ for all $i \in I$. Let $\mathcal{B}(\lambda)$ denote the crystal basis of the extremal weight module $V(\lambda)$ over $U_q(\mathfrak{g}_{\text{af}})$, and let $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ denote the set of SiLS paths of shape λ , equipped with the $U_q(\mathfrak{g}_{\text{af}})$ -crystal structure as above. Then, we have an isomorphism of crystals*

$$\mathcal{B}(\lambda) \cong \mathbb{B}^{\frac{\infty}{2}}(\lambda). \quad (2)$$

Remark that for each $i \in I$, we have a natural identification $\mathbb{B}^{\frac{\infty}{2}}(m_i \varpi_i) = \mathcal{B}(m_i \varpi_i)$, since the equality $(W_{I \setminus \{i\}})_{\text{af}} = (W_{\text{af}})_{m_i \varpi_i}$ holds. Hence we recover our previous results in [NS2, NS3]. Also, we should mention that Hernandez and Nakajima ([HN]) gave a monomial realization of the connected component $\mathcal{B}_0(\lambda)$; however, their realization is given in a recursive way, and it is difficult to determine all the elements in $\mathcal{B}_0(\lambda)$ explicitly in this realization.

As an application of the isomorphism theorem above, in our paper [NS5] sequel to the present one, we prove that the graded character of a particular Demazure submodule $V_{w_0}^+(\lambda)$ of $V(\lambda)$

for the longest element $w_0 \in W$ is identical to a q -Whittaker function, which is nothing but the graded character of a global Weyl module over the current algebra (see [BF] for simply-laced cases). This result yields a purely crystal-theoretical explanation of the relation between a q -Whittaker function and the specialization at $t = 0$ of a symmetric Macdonald polynomial, which is nothing but the graded character of a local Weyl module over the current algebra (see [LNS³2]).

This paper is organized as follows. In §2, we fix our notation for untwisted affine root data, and review some basic facts about LS paths and (parabolic) semi-infinite Bruhat graphs. In §3, we introduce SiLS paths and define a crystal structure on them, postponing to §4 the proof of the stability property of the set of semi-infinite LS paths under root operators. Also, we state our main result, i.e., the isomorphism theorem above between $\mathcal{B}(\lambda)$ and $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$. In §4, we first study some fundamental properties of semi-infinite Bruhat graphs, and then prove that there exists a canonical surjection from $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ to $\mathbb{B}(\lambda)$. Using this surjection, we give the postponed proof of the stability property above of SiLS paths. In §5, we prove that the connected component $\mathcal{B}_0(\lambda)$ of $\mathcal{B}(\lambda)$ containing u_λ is isomorphic, as a crystal, to the connected component $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ containing the element $\eta_e := (e; 0, 1)$. In §6, we obtain a condition for the existence of a directed path in a (parabolic) semi-infinite Bruhat graph from a translation to another translation; this result is used in §7 to give a parametrization of the connected components of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$. Finally, in §7, by combining the results in §4, §5, and §6, we obtain the desired isomorphism $\mathcal{B}(\lambda) \cong \mathbb{B}^{\frac{\infty}{2}}(\lambda)$. In Appendix A, we give another (but equivalent) definition of semi-infinite Bruhat graphs, and also mention the relation between the semi-infinite Bruhat order and Lusztig's generic Bruhat order.

Acknowledgements. The second and third authors would like to thank Christian Lenart, Anne Schilling, and Mark Shimozono for related collaborations. The third author thanks Professor Peter Littelmann, Professor Ghislain Fourier, Dr. Deniz Kus, and students attending at Professor Littelmann's seminar for giving us valuable comments on this work.

2 LS paths and semi-infinite Bruhat graphs.

2.1 Untwisted affine root data.

Let $\mathfrak{g}_{\text{af}}$ be an untwisted affine Lie algebra over $\mathbb{C}$ with Cartan subalgebra $\mathfrak{h}_{\text{af}}$. Let $\{\alpha_i\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}^* := \text{Hom}_{\mathbb{C}}(\mathfrak{h}_{\text{af}}, \mathbb{C})$ and $\{\alpha_i^\vee\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}$ be the sets of simple roots and simple coroots, respectively. Let $\langle \cdot, \cdot \rangle : \mathfrak{h}_{\text{af}} \times \mathfrak{h}_{\text{af}}^* \rightarrow \mathbb{C}$ denote the canonical pairing. Throughout this paper, we take and fix an integral weight lattice $P_{\text{af}} \subset \mathfrak{h}_{\text{af}}^*$ satisfying the conditions that $\alpha_i \in P_{\text{af}}$ and $\alpha_i^\vee \in \text{Hom}_{\mathbb{Z}}(P_{\text{af}}, \mathbb{Z})$ for all $i \in I_{\text{af}}$, and that for each $i \in I_{\text{af}}$ there exists $\Lambda_i \in P_{\text{af}}$ such that $\langle \alpha_j^\vee, \Lambda_i \rangle = \delta_{ij}$ for all $j \in I_{\text{af}}$. Let $\delta = \sum_{i \in I_{\text{af}}} a_i \alpha_i \in \mathfrak{h}_{\text{af}}^*$ and $c = \sum_{i \in I_{\text{af}}} a_i^\vee \alpha_i^\vee \in \mathfrak{h}_{\text{af}}$ be the null root

and the canonical central element, respectively. We take and fix $0 \in I_{\text{af}}$ such that $a_0 = a_0^\vee = 1$, and set $I := I_{\text{af}} \setminus \{0\}$; note that the subset I of I_{af} corresponds to the index set for the finite-dimensional simple Lie subalgebra $\mathfrak{g}$ of $\mathfrak{g}_{\text{af}}$. For each $i \in I$, we define $\varpi_i := \Lambda_i - \langle c, \Lambda_i \rangle \Lambda_0$; note that $\langle c, \varpi_i \rangle = 0$ for all $i \in I$. Set

$$Q := \bigoplus_{i \in I} \mathbb{Z} \alpha_i, \quad Q^\vee := \bigoplus_{i \in I} \mathbb{Z} \alpha_i^\vee, \quad P^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0} \varpi_i;$$

we call an element of P^+ a level-zero dominant integral weight.

Let $W_{\text{af}} := \langle r_i \mid i \in I_{\text{af}} \rangle$ be the (affine) Weyl group of $\mathfrak{g}_{\text{af}}$, where r_i denotes the simple reflection with respect to α_i , and set $W := \langle r_i \mid i \in I \rangle \subset W_{\text{af}}$, which can be regarded as the Weyl group of $\mathfrak{g}$. Let $e \in W_{\text{af}}$ be the unit element, and $\ell : W_{\text{af}} \rightarrow \mathbb{Z}_{\geq 0}$ the length function. For $\xi \in Q^\vee$, denote by $t_\xi \in W_{\text{af}}$ the translation with respect to ξ (see [Kac, §6.5]). We know from [Kac, Proposition 6.5] that $\{t_\xi \mid \xi \in Q^\vee\}$ forms an abelian normal subgroup of W_{af} , for which $t_\xi t_\zeta = t_{\xi+\zeta}$, $\xi, \zeta \in Q^\vee$, and $W_{\text{af}} = W \ltimes \{t_\xi \mid \xi \in Q^\vee\}$; remark that for $w \in W$ and $\xi \in Q^\vee$, we have

$$wt_\xi \mu = w\mu - \langle \xi, \mu \rangle \delta \quad \text{if } \mu \in \mathfrak{h}_{\text{af}}^* \text{ satisfies } \langle c, \mu \rangle = 0. \quad (2.1.1)$$

Denote by Δ_{af} the set of real roots of $\mathfrak{g}_{\text{af}}$, and Δ_{af}^+ the set of positive real roots of $\mathfrak{g}_{\text{af}}$; we know from [Kac, Proposition 6.3] that

$$\begin{aligned} \Delta_{\text{af}} &= \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}\}, \\ \Delta_{\text{af}}^+ &= \Delta^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}_{>0}\}, \end{aligned}$$

where $\Delta := \Delta_{\text{af}} \cap Q$ is the (finite) root system corresponding to I , and $\Delta^+ := \Delta \cap \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i$. For $\beta \in \Delta_{\text{af}}$, denote by $\beta^\vee \in Q^\vee$ the coroot of β , and by $r_\beta \in W_{\text{af}}$ the reflection with respect to β ; if $\beta \in \Delta_{\text{af}}$ is of the form $\beta = \alpha + n\delta$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}$, then

$$r_\beta = r_\alpha t_{n\alpha^\vee}. \quad (2.1.2)$$

For a subset J of I , we set

$$\begin{aligned} Q_J &:= \bigoplus_{j \in J} \mathbb{Z} \alpha_j, & Q_J^\vee &:= \bigoplus_{j \in J} \mathbb{Z} \alpha_j^\vee, & Q_J^{\vee+} &:= \sum_{j \in J} \mathbb{Z}_{\geq 0} \alpha_j^\vee, \\ \Delta_J &:= \Delta \cap Q_J, & \Delta_J^+ &:= \Delta_J \cap \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i, & W_J &:= \langle r_j \mid j \in J \rangle. \end{aligned}$$

Let W^J denote the set of minimal(-length) coset representatives for W/W_J ; we see from [BB, §2.4] that

$$W^J = \{w \in W \mid w\alpha \in \Delta^+ \text{ for all } \alpha \in \Delta_J^+\}. \quad (2.1.3)$$

For $w \in W$, we denote by $[w] = [w]^J \in W^J$ the minimal coset representative for the coset wW_J in W/W_J .

2.2 Lakshmibai-Seshadri paths.

In this subsection, we briefly review some basic facts about crystals of Lakshmibai-Seshadri (LS for short) paths, introduced by Littelmann [Li1], [Li2]. In this subsection, we fix $\lambda \in P^+$.

Definition 2.2.1 ([Li2, §4]). We define a partial order $\leq$ on $W_{\text{af}}\lambda$ as follows: for $\mu, \nu \in W_{\text{af}}\lambda$, we write $\mu \leq \nu$ if there exists a sequence $\mu = \nu_0, \nu_1, \dots, \nu_k = \nu$ of elements in $W_{\text{af}}\lambda$ and a sequence $\beta_1, \dots, \beta_k$ of elements in Δ_{af}^+ such that $\nu_m = r_{\beta_m} \nu_{m-1}$ and $\langle \beta_m^\vee, \nu_{m-1} \rangle \in \mathbb{Z}_{>0}$ for all $m = 1, 2, \dots, k$. We call the poset $(W_{\text{af}}\lambda, \leq)$ the level-zero weight poset of shape λ .

Remark 2.2.2. Let $\nu \in P$, and $\beta, \gamma \in \Delta_{\text{af}}^+$. If $r_\beta \nu = r_\gamma \nu \neq \nu$, then $\beta = \gamma$. Indeed, we have $\nu - \langle \beta^\vee, \nu \rangle \beta = \nu - \langle \gamma^\vee, \nu \rangle \gamma$, and hence $\langle \beta^\vee, \nu \rangle \beta = \langle \gamma^\vee, \nu \rangle \gamma$. Also, since $r_\beta \nu \neq \nu$ and $r_\gamma \nu \neq \nu$, it follows immediately that $\langle \beta^\vee, \nu \rangle \neq 0$ and $\langle \gamma^\vee, \nu \rangle \neq 0$. Hence, by [Kac, Proposition 5.1 b)], we obtain $\beta = \gamma$.

Definition 2.2.3.

- (1) Define $\text{LZ}(\lambda)$ to be the Δ_{af}^+ -labeled, directed graph with vertex set $W_{\text{af}}\lambda$ and Δ_{af}^+ -labeled, directed edges of the following form: $\mu \xrightarrow{\beta} \nu$ for $\mu, \nu \in W_{\text{af}}\lambda$, where ν covers μ in the poset $W_{\text{af}}\lambda$, and the label β of the edge is a unique positive real root $\beta \in \Delta_{\text{af}}^+$ such that $\nu = r_\beta \mu$ and $\langle \beta^\vee, \mu \rangle > 0$ (see Remark 2.2.2).
- (2) Let $0 < a \leq 1$ be a rational number. Define $\text{LZ}(\lambda; a)$ to be the subgraph of $\text{LZ}(\lambda)$ with the same vertex set but having only the edges of the form:

$$\mu \xrightarrow{\beta} \nu \quad \text{with} \quad a \langle \beta^\vee, \mu \rangle \in \mathbb{Z}; \quad (2.2.1)$$

note that $\text{LZ}(\lambda; 1) = \text{LZ}(\lambda)$.

Definition 2.2.4 ([Li2, §4]). An LS path of shape λ is, by definition, a pair $(\boldsymbol{\nu}; \mathbf{a})$ of a decreasing sequence $\boldsymbol{\nu} : \nu_1 > \dots > \nu_s$ of elements in $W_{\text{af}}\lambda$ and an increasing sequence $\mathbf{a} : 0 = a_0 < a_1 < \dots < a_s = 1$ of rational numbers satisfying the condition that there exists a directed path from ν_{u+1} to ν_u in $\text{LZ}(\lambda; a_u)$ for each $u = 1, 2, \dots, s-1$. Let $\mathbb{B}(\lambda)$ denote the set of LS paths of shape λ .

We identify $\pi = (\nu_1, \dots, \nu_s; a_0, \dots, a_s) \in \mathbb{B}(\lambda)$ with the piecewise-linear, continuous map $\pi : [0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} P_{\text{af}}$ whose “direction vector” for the interval $[a_{u-1}, a_u]$ is equal to ν_u for each $1 \leq u \leq s$, that is,

$$\pi(t) = \sum_{p=1}^{u-1} (a_p - a_{p-1}) \nu_p + (t - a_{u-1}) \nu_u \quad \text{for } t \in [a_{u-1}, a_u], \quad 1 \leq u \leq s. \quad (2.2.2)$$

We also express $\pi = (\nu_1, \dots, \nu_s; a_0, \dots, a_s) \in \mathbb{B}(\lambda)$ as:

$$0 = a_0 \xrightarrow{\nu_1} a_1 \xrightarrow{\nu_2} \dots \xrightarrow{\nu_{s-1}} a_{s-1} \xrightarrow{\nu_s} a_s = 1.$$

Now, we equip the set $\mathbb{B}(\lambda)$ with a crystal structure with weights in P_{af} as follows (for the definition of crystals, see [Kas2, §7.2] and [HK, Definition 4.5.1] for example). First, we define $\text{wt} : \mathbb{B}(\lambda) \rightarrow P_{\text{af}}$ by $\text{wt}(\pi) := \pi(1) \in P_{\text{af}}$; we know from [Li2, Lemma 4.5 a)] that $\pi(1) \in P_{\text{af}}$ for all $\pi \in \mathbb{B}(\lambda)$. Next, for $\pi \in \mathbb{B}(\lambda)$ and $i \in I_{\text{af}}$, we set

$$\begin{cases} H_i^\pi(t) := \langle \alpha_i^\vee, \pi(t) \rangle & \text{for } t \in [0, 1], \\ m_i^\pi := \min\{H_i^\pi(t) \mid t \in [0, 1]\}. \end{cases} \quad (2.2.3)$$

Remark 2.2.5. We see from [Li2, Lemma 4.5 d)] that for each $\pi \in \mathbb{B}(\lambda)$ and $i \in I_{\text{af}}$, all local minima of the function $H_i^\pi(t)$, $t \in [0, 1]$, are integers. In particular, the minimum m_i^π is a nonpositive integer (recall that $\pi(0) = 0$, and hence $H_i^\pi(0) = 0$).

Following [Li2, §1] (see also [NS3, §1]), we define the root operators e_i, f_i , $i \in I_{\text{af}}$, on $\mathbb{B}(\lambda) \sqcup \{\mathbf{0}\}$ as follows. Here, $\mathbf{0}$ is an additional element not contained in any crystal.

Definition 2.2.6. Let $\pi = (\nu_1, \dots, \nu_s; a_0, \dots, a_s) \in \mathbb{B}(\lambda)$, and $i \in I_{\text{af}}$.

(1) If $m_i^\pi = 0$, then we define $e_i\pi := \mathbf{0}$. If $m_i^\pi \leq -1$, then set

$$\begin{cases} t_1 := \min\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_0 := \max\{t \in [0, t_1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (2.2.4)$$

we deduce from Remark 2.2.5 that $H_i^\pi(t)$ is strictly decreasing on $[t_0, t_1]$. Notice that there exists $1 \leq q \leq s$ such that $t_1 = a_q$. Let $1 \leq p \leq q$ be such that $a_{p-1} \leq t_0 < a_p$. Then we define $e_i\pi$ to be

$$0 = a_0 \xrightarrow{\nu_1} \dots \xrightarrow{\nu_{p-1}} a_{p-1} \xrightarrow{\nu_p} \underbrace{t_0 \xrightarrow{r_i\nu_p} a_p \xrightarrow{r_i\nu_{p+1}} \dots \xrightarrow{r_i\nu_q} a_q}_{\text{"reflected" by } r_i} = t_1 \xrightarrow{\nu_{q+1}} \dots \xrightarrow{\nu_s} a_s = 1,$$

that is,

$$\begin{aligned} e_i\pi := & (\nu_1, \dots, \nu_p, r_i\nu_p, r_i\nu_{p+1}, \dots, r_i\nu_q, \nu_{q+1}, \dots, \nu_s; \\ & a_0, \dots, a_{p-1}, t_0, a_p, \dots, a_q = t_1, \dots, a_s); \end{aligned}$$

if $t_0 = a_{p-1}$, then we drop ν_p and a_{p-1} , and if $r_i\nu_q = \nu_{q+1}$, then we drop ν_{q+1} and $a_q = t_1$.

(2) If $H_i^\pi(1) - m_i^\pi = 0$, then we define $f_i\pi := \mathbf{0}$. If $H_i^\pi(1) - m_i^\pi \geq 1$, then set

$$\begin{cases} t_0 := \max\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_1 := \min\{t \in [t_0, 1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (2.2.5)$$

we deduce from Remark 2.2.5 that $H_i^\pi(t)$ is strictly increasing on $[t_0, t_1]$. Notice that there exists $0 \leq p \leq s-1$ such that $t_0 = a_p$. Let $p \leq q \leq s-1$ be such that $a_q < t_1 \leq a_{q+1}$. Then we define $f_i\pi$ to be

$$0 = a_0 \xrightarrow{\nu_1} \dots \xrightarrow{\nu_p} a_p = t_0 \xrightarrow{r_i\nu_{p+1}} \dots \xrightarrow{r_i\nu_q} a_q \xrightarrow{r_i\nu_{q+1}} t_1 \xrightarrow{\nu_{q+1}} a_{q+1} \xrightarrow{\nu_{q+2}} \dots \xrightarrow{\nu_s} a_s = 1,$$

"reflected" by r_i

that is,

$$\begin{aligned} f_i \pi &:= (\nu_1, \dots, \nu_p, r_i \nu_{p+1}, \dots, r_i \nu_q, r_i \nu_{q+1}, \nu_{q+1}, \dots, \nu_s; \\ &\quad a_0, \dots, a_p = t_0, \dots, a_q, t_1, a_{q+1}, \dots, a_s); \end{aligned}$$

if $t_1 = a_{q+1}$, then we drop ν_{q+1} and a_{q+1} , and if $\nu_p = r_i \nu_{p+1}$, then we drop ν_p and $a_p = t_0$.

(3) Set $e_i \mathbf{0} = f_i \mathbf{0} := \mathbf{0}$ for all $i \in I_{\text{af}}$.

We know from [Li2, Corollary 2 a)] that the set $\mathbb{B}(\lambda) \sqcup \{\mathbf{0}\}$ is stable under the action of the root operators $e_i, f_i, i \in I_{\text{af}}$. Now we define

$$\begin{cases} \varepsilon_i(\pi) := \max\{n \in \mathbb{Z}_{\geq 0} \mid e_i^n \pi \neq \mathbf{0}\}, \\ \varphi_i(\pi) := \max\{n \in \mathbb{Z}_{\geq 0} \mid f_i^n \pi \neq \mathbf{0}\} \end{cases} \quad (2.2.6)$$

for $\pi \in \mathbb{B}(\lambda)$ and $i \in I_{\text{af}}$. We know from [Li2, §2 and §4] that the set $\mathbb{B}(\lambda)$, equipped with the maps $\text{wt}, e_i, f_i, i \in I_{\text{af}}$, and $\varepsilon_i, \varphi_i, i \in I$, defined above, is a crystal with weights in P_{af} .

2.3 Peterson's coset representatives.

Let J be a subset of I . Following [P] (see also [LS, §10]), we define

$$(\Delta_J)_{\text{af}} := \{\alpha + n\delta \mid \alpha \in \Delta_J, n \in \mathbb{Z}\} \subset \Delta_{\text{af}}, \quad (2.3.1)$$

$$(\Delta_J)_{\text{af}}^+ := (\Delta_J)_{\text{af}} \cap \Delta_{\text{af}}^+ = \Delta_J^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta_J, n \in \mathbb{Z}_{>0}\}, \quad (2.3.2)$$

$$(W_J)_{\text{af}} := W_J \ltimes \{t_\xi \mid \xi \in Q_J^\vee\}, \quad (2.3.3)$$

$$(W^J)_{\text{af}} := \{x \in W_{\text{af}} \mid x\beta \in \Delta_{\text{af}}^+ \text{ for all } \beta \in (\Delta_J)_{\text{af}}^+\}. \quad (2.3.4)$$

Remark 2.3.1. We can easily show, using (2.1.2), that $(W_J)_{\text{af}} = \langle r_\beta \mid \beta \in (\Delta_J)_{\text{af}}^+ \rangle$.

We know the following proposition from [P] (see also [LS, Lemma 10.6]).

Proposition 2.3.2. *For each $x \in W_{\text{af}}$, there exist a unique $x_1 \in (W^J)_{\text{af}}$ and a unique $x_2 \in (W_J)_{\text{af}}$ such that $x = x_1 x_2$.*

We define a (surjective) map $\Pi^J : W_{\text{af}} \rightarrow (W^J)_{\text{af}}$ by $\Pi^J(x) := x_1$ if $x = x_1 x_2$ with $x_1 \in (W^J)_{\text{af}}$ and $x_2 \in (W_J)_{\text{af}}$.

Lemma 2.3.3 ([P]; see also [LS, Proposition 10.10]).

(1) $\Pi^J(w) = \lfloor w \rfloor$ for every $w \in W$.

(2) $\Pi^J(x t_\xi) = \Pi^J(x) \Pi^J(t_\xi)$ for every $x \in W_{\text{af}}$ and $\xi \in Q^\vee$.

Definition 2.3.4 (see [LNS³1, Lemma 3.8]). An element $\xi \in Q^\vee$ is said to be J -adjusted if $\langle \xi, \gamma \rangle \in \{-1, 0\}$ for all $\gamma \in \Delta_J^+$. Let $Q^{\vee, J\text{-ad}}$ denote the set of J -adjusted elements.

Lemma 2.3.5.

- (1) For each $\xi \in Q^\vee$, there exists a unique $\phi_J(\xi) \in Q_J^\vee$ such that $\xi + \phi_J(\xi) \in Q^{\vee, J\text{-ad}}$. In particular, $\xi \in Q^{\vee, J\text{-ad}}$ if and only if $\phi_J(\xi) = 0$.
- (2) For each $\xi \in Q^\vee$, the element $\Pi^J(t_\xi) \in (W^J)_{\text{af}}$ is of the form $\Pi^J(t_\xi) = z_\xi t_{\xi + \phi_J(\xi)}$ for some $z_\xi \in W_J$. Therefore, by Lemma 2.3.3, $\Pi^J(wt_\xi) = [w]z_\xi t_{\xi + \phi_J(\xi)}$ for every $w \in W$ and $\xi \in Q^\vee$.
- (3) We have

$$(W^J)_{\text{af}} = \{wz_\xi t_\xi \mid w \in W^J, \xi \in Q^{\vee, J\text{-ad}}\}. \quad (2.3.5)$$

Proof. Parts (1), (2), and (3) follow immediately from [LNS³1, (3.6), (3.7), and Lemma 3.7], respectively. $\square$

Now we prove a few easy lemmas, which will be used later.

Lemma 2.3.6. Let $x \in (W^J)_{\text{af}}$ and $i \in I_{\text{af}}$. Then, $x^{-1}\alpha_i \notin (\Delta_J)_{\text{af}}$ if and only if $r_i x \in (W^J)_{\text{af}}$.

Proof. First, let us show the “only if” part; by definition (2.3.4) of $(W^J)_{\text{af}}$, it suffices to show that $r_i x \beta \in \Delta_{\text{af}}^+$ for all $\beta \in (\Delta_J)_{\text{af}}^+$. Let $\beta \in (\Delta_J)_{\text{af}}^+$. Since $x \in (W^J)_{\text{af}}$, we have $x\beta \in \Delta_{\text{af}}^+$. Also, since $x^{-1}\alpha_i \notin (\Delta_J)_{\text{af}}$ by the assumption, it follows that $x^{-1}\alpha_i \neq \beta$, and hence $x\beta \neq \alpha_i$. Therefore, we obtain $r_i x \beta \in \Delta_{\text{af}}^+$.

Next, let us show the “if” part. Suppose, for a contradiction, that $x^{-1}\alpha_i \in (\Delta_J)_{\text{af}}$. Then, since $r_{x^{-1}\alpha_i} \in (W_J)_{\text{af}}$, we see by Proposition 2.3.2 that $\Pi^J(r_i x) = \Pi^J(xr_{x^{-1}\alpha_i}) = \Pi^J(x)$. Since $r_i x, x \in (W^J)_{\text{af}}$ by the assumption, we have $\Pi^J(r_i x) = r_i x$ and $\Pi^J(x) = x$. Therefore, we obtain $r_i x = x$, which is a contradiction. Thus, $x^{-1}\alpha_i \notin (\Delta_J)_{\text{af}}$. This proves the lemma. $\square$

Lemma 2.3.7. Let $x \in W_{\text{af}}$, and $\xi \in Q^{\vee, J\text{-ad}}$. Then, $xz_\xi t_\xi \in (W^J)_{\text{af}}$ if and only if $x \in (W^J)_{\text{af}}$.

Proof. First, we remark that

$$\begin{aligned} \Pi^J(xz_\xi t_\xi) &= \Pi^J(xz_\xi)\Pi^J(t_\xi) && \text{by Lemma 2.3.3 (2)} \\ &= \Pi^J(x)z_\xi t_\xi && \text{by Lemmas 2.3.3 (1) and 2.3.5 (1), (2).} \end{aligned} \quad (2.3.6)$$

Let us prove the “only if” part. Assume that $xz_\xi t_\xi \in (W^J)_{\text{af}}$; note that $\Pi^J(xz_\xi t_\xi) = xz_\xi t_\xi$. Combining this equality and (2.3.6), we obtain $\Pi^J(x)z_\xi t_\xi = xz_\xi t_\xi$, and hence $\Pi^J(x) = x$, which implies that $x \in (W^J)_{\text{af}}$. Now, let us prove the “if” part. Assume that $x \in (W^J)_{\text{af}}$; note that $\Pi^J(x) = x$. Combining this equality and (2.3.6), we obtain $\Pi^J(xz_\xi t_\xi) = \Pi^J(x)z_\xi t_\xi = xz_\xi t_\xi$, which implies that $xz_\xi t_\xi \in (W^J)_{\text{af}}$. This proves the lemma. $\square$

2.4 Semi-infinite Bruhat graphs.

Definition 2.4.1 ([P]). Let $x \in W_{\text{af}}$, and write it as $x = vt_\zeta$ with $v \in W$ and $\zeta \in Q^\vee$. Then we define the semi-infinite length $\ell^{\frac{\infty}{2}}(x)$ of x by

$$\ell^{\frac{\infty}{2}}(x) := \ell(v) + 2\langle \zeta, \rho \rangle,$$

where $\rho := (1/2) \sum_{\alpha \in \Delta^+} \alpha$.

Definition 2.4.2. Let J be a subset of I .

- (1) Define the (parabolic) semi-infinite Bruhat graph SiB^J to be the Δ_{af}^+ -labeled, directed graph with vertex set $(W^J)_{\text{af}}$ and Δ_{af}^+ -labeled, directed edges of the following form: $x \xrightarrow{\beta} r_\beta x$ for $x \in (W^J)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$, where $r_\beta x \in (W^J)_{\text{af}}$ and $\ell^{\frac{\infty}{2}}(r_\beta x) = \ell^{\frac{\infty}{2}}(x) + 1$.
- (2) The semi-infinite Bruhat order is a partial order $\leq_{\frac{\infty}{2}}$ on $(W^J)_{\text{af}}$ defined as follows: for $x, y \in (W^J)_{\text{af}}$, we write $x \leq_{\frac{\infty}{2}} y$ if there exists a directed path from x to y in SiB^J .

3 Isomorphism theorem.

3.1 Semi-infinite Lakshmibai-Seshadri paths.

Definition 3.1.1. Let $\lambda \in P^+$, and set $J = J_\lambda := \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\} \subset I$. For a rational number $0 < a \leq 1$, define $\text{SiB}(\lambda; a)$ to be the subgraph of SiB^J with the same vertex set but having only the edges of the form:

$$x \xrightarrow{\beta} y \quad \text{with} \quad a\langle \beta^\vee, x\lambda \rangle \in \mathbb{Z}; \quad (3.1.1)$$

note that $\text{SiB}(\lambda; 1) = \text{SiB}^J$.

Definition 3.1.2. Let $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. A semi-infinite Lakshmibai-Seshadri (SiLS for short) path of shape λ is, by definition, a pair $(\mathbf{x}; \mathbf{a})$ of a decreasing sequence $\mathbf{x} : x_1 >_{\frac{\infty}{2}} \cdots >_{\frac{\infty}{2}} x_s$ of elements in $(W^J)_{\text{af}}$ and an increasing sequence $\mathbf{a} : 0 = a_0 < a_1 < \cdots < a_s = 1$ of rational numbers satisfying the condition that there exists a directed path from x_{u+1} to x_u in $\text{SiB}(\lambda; a_u)$ for each $u = 1, 2, \dots, s-1$; we express this element as:

$$0 = a_0 \xrightarrow{x_1} a_1 \xrightarrow{x_2} \cdots \xrightarrow{x_{s-1}} a_{s-1} \xrightarrow{x_s} a_s = 1.$$

Denote by $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ the set of all SiLS paths of shape λ .

For the rest of this subsection, we fix $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. For $\eta = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, we set

$$\overline{\eta} := (x_1\lambda, \dots, x_s\lambda; a_0, \dots, a_s).$$

Proposition 3.1.3 (which will be proved in §4.2). *It holds that $\bar{\eta} \in \mathbb{B}(\lambda)$ for every $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$. Thus we obtain a map $^- : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \rightarrow \mathbb{B}(\lambda)$, $\eta \mapsto \bar{\eta}$.*

We equip the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ with a crystal structure with weights in P_{af} as follows. We define $\text{wt} : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \rightarrow P_{\text{af}}$ by: $\text{wt}(\eta) := \text{wt}(\bar{\eta}) = \bar{\eta}(1) \in P_{\text{af}}$. We define operators $e_i, f_i, i \in I_{\text{af}}$, which we call root operators for $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$, in the same manner as for $\mathbb{B}(\lambda)$.

Definition 3.1.4. Let $\eta = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, and $i \in I_{\text{af}}$.

- (1) If $m_i^{\bar{\eta}} = 0$, then we define $e_i \eta := \mathbf{0}$. If $m_i^{\bar{\eta}} \leq -1$, then define $t_0, t_1 \in [0, 1]$ by (2.2.4), with $\pi = \bar{\eta}$. Let $1 \leq p \leq q \leq s$ be such that $a_{p-1} \leq t_0 < a_p$ and $t_1 = a_q$. Then we define $e_i \eta$ to be

$$0 = a_0 \xrightarrow{x_1} \dots \xrightarrow{x_{p-1}} a_{p-1} \xrightarrow{x_p} \underbrace{t_0 \xrightarrow{r_i x_p} a_p \xrightarrow{r_i x_{p+1}} \dots \xrightarrow{r_i x_q} a_q}_{\text{"reflected" by } r_i} = t_1 \xrightarrow{x_{q+1}} \dots \xrightarrow{x_s} a_s = 1,$$

that is,

$$e_i \eta := (x_1, \dots, x_p, r_i x_p, r_i x_{p+1}, \dots, r_i x_q, x_{q+1}, \dots, x_s; \\ a_0, \dots, a_{p-1}, t_0, a_p, \dots, a_q = t_1, \dots, a_s);$$

if $t_0 = a_{p-1}$, then we drop x_p and a_{p-1} , and if $r_i x_q = x_{q+1}$, then we drop x_{q+1} and $a_q = t_1$.

- (2) If $H_i^{\bar{\eta}}(1) - m_i^{\bar{\eta}} = 0$, then we define $f_i \eta := \mathbf{0}$. If $H_i^{\bar{\eta}}(1) - m_i^{\bar{\eta}} \geq 1$, then define $t_0, t_1 \in [0, 1]$ by (2.2.5), with $\pi = \bar{\eta}$. Let $0 \leq p \leq q \leq s-1$ be such that $t_0 = a_p$, and $a_q < t_1 \leq a_{q+1}$. Then we define $f_i \eta$ to be

$$0 = a_0 \xrightarrow{x_1} \dots \xrightarrow{x_p} a_p = t_0 \xrightarrow{r_i x_{p+1}} \dots \xrightarrow{r_i x_q} a_q \xrightarrow{r_i x_{q+1}} t_1 \xrightarrow{x_{q+1}} a_{q+1} \xrightarrow{x_{q+2}} \dots \xrightarrow{x_s} a_s = 1,$$

"reflected" by r_i

that is,

$$f_i \eta := (x_1, \dots, x_p, r_i x_{p+1}, \dots, r_i x_q, r_i x_{q+1}, x_{q+1}, \dots, x_s; \\ a_0, \dots, a_p = t_0, \dots, a_q, t_1, a_{q+1}, \dots, a_s);$$

if $t_1 = a_{q+1}$, then we drop x_{q+1} and a_{q+1} , and if $x_p = r_i x_{p+1}$, then we drop x_p and $a_p = t_0$.

- (3) Set $e_i \mathbf{0} = f_i \mathbf{0} := \mathbf{0}$ for all $i \in I_{\text{af}}$.

Theorem 3.1.5 (which will be proved in §4.3).

- (1) *The set $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \sqcup \{\mathbf{0}\}$ is stable under the action of the root operators e_i and $f_i, i \in I_{\text{af}}$.*
- (2) *For each $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$, we set*

$$\begin{cases} \varepsilon_i(\eta) := \max\{n \geq 0 \mid e_i^n \eta \neq \mathbf{0}\}, \\ \varphi_i(\eta) := \max\{n \geq 0 \mid f_i^n \eta \neq \mathbf{0}\}. \end{cases}$$

Then, the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$, equipped with the maps $\text{wt}, e_i, f_i, i \in I_{\text{af}}$, and $\varepsilon_i, \varphi_i, i \in I_{\text{af}}$, defined above, is a crystal with weights in P_{af} .

3.2 Isomorphism theorem between $\mathcal{B}(\lambda)$ and $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$.

Let $V(\lambda)$ denote the extremal weight module of extremal weight $\lambda \in P^+$ over the quantized universal enveloping algebra $U_q(\mathfrak{g}_{\text{af}})$ associated with $\mathfrak{g}_{\text{af}}$, which is an integrable $U_q(\mathfrak{g}_{\text{af}})$ -module generated by a single element v_λ with the defining relation that v_λ is an “extremal weight vector” of weight λ (for details, see [Kas1, §8] and [Kas5, §3]). We know from [Kas1, §8] that $V(\lambda)$ has a crystal basis $\mathcal{B}(\lambda)$. The main result of this paper is the following theorem.

Theorem 3.2.1. *Let $\lambda \in P^+$. The crystal basis $\mathcal{B}(\lambda)$ of the extremal weight module $V(\lambda)$ of extremal weight λ is isomorphic, as a crystal, to the crystal $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ of SiLS paths of shape λ .*

Let us give an outline of the proof of Theorem 3.2.1. Let $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ denote the connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ containing $\eta_e := (e; 0, 1) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$. Also, let u_λ be the element of $\mathcal{B}(\lambda)$ corresponding to the generator v_λ of $V(\lambda)$, and let $\mathcal{B}_0(\lambda)$ denote the connected component of $\mathcal{B}(\lambda)$ containing $u_\lambda \in \mathcal{B}(\lambda)$.

Proposition 3.2.2 (which will be proved in §5). *There exists a unique isomorphism $\mathcal{B}_0(\lambda) \xrightarrow{\sim} \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals that maps u_λ to η_e .*

We write $\lambda \in P^+$ as $\lambda = \sum_{i \in I} m_i \varpi_i$ with $m_i \in \mathbb{Z}_{\geq 0}$, $i \in I$, and define

$$\text{Par}(\lambda) := \{ \boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \mid \rho^{(i)} \text{ is a partition of length less than } m_i \text{ for each } i \in I \}; \quad (3.2.1)$$

we understand that $\rho^{(i)}$ is the empty partition if $m_i = 0$. We equip the set $\text{Par}(\lambda)$ with a crystal structure as follows: for each $\boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \in \text{Par}(\lambda)$, we set

$$\begin{cases} e_i \boldsymbol{\rho} = f_i \boldsymbol{\rho} := \mathbf{0}, \quad \varepsilon_i(\boldsymbol{\rho}) = \varphi_i(\boldsymbol{\rho}) := -\infty & \text{for } i \in I_{\text{af}}, \\ \text{wt}(\boldsymbol{\rho}) := -\sum_{i \in I} |\rho^{(i)}| \delta, \end{cases} \quad (3.2.2)$$

where for a partition $\chi = (\chi_1 \geq \chi_2 \geq \cdots \geq \chi_k \geq 0)$, we set $|\chi| := \sum_{l=1}^k \chi_l$. By Proposition 3.2.2, we have the isomorphism

$$\text{Par}(\lambda) \otimes \mathcal{B}_0(\lambda) \cong \text{Par}(\lambda) \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda) \quad (3.2.3)$$

of crystals. Let $\mathcal{B}$ be either $\mathcal{B}_0(\lambda)$ or $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$. For each $\boldsymbol{\rho} \in \text{Par}(\lambda)$, we set $\{\boldsymbol{\rho}\} \otimes \mathcal{B} := \{ \boldsymbol{\rho} \otimes b \mid b \in \mathcal{B} \} \subset \text{Par}(\lambda) \otimes \mathcal{B}$. Then it is easily seen from the tensor product rule for crystals that

$$\text{Par}(\lambda) \otimes \mathcal{B} = \bigsqcup_{\boldsymbol{\rho} \in \text{Par}(\lambda)} \{\boldsymbol{\rho}\} \otimes \mathcal{B}$$

is the decomposition of $\text{Par}(\lambda) \otimes \mathcal{B}$ into its connected components. Moreover, the map $\mathcal{B} \rightarrow \{\boldsymbol{\rho}\} \otimes \mathcal{B}$, $b \mapsto \boldsymbol{\rho} \otimes b$, is bijective and commutes with Kashiwara operators.

Now, we know the following proposition from [BN, Theorem 4.16 (i)].

Proposition 3.2.3. *For $\lambda \in P^+$, there exists an isomorphism $\mathcal{B}(\lambda) \xrightarrow{\sim} \text{Par}(\lambda) \otimes \mathcal{B}_0(\lambda)$ of crystals.*

Also, we have the following proposition.

Proposition 3.2.4 (which will be proved in §7.2). *For $\lambda \in P^+$, there exists an isomorphism $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \xrightarrow{\sim} \text{Par}(\lambda) \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals.*

Combining all the results above, we finally obtain

$$\begin{aligned} \mathcal{B}(\lambda) &\cong \text{Par}(\lambda) \otimes \mathcal{B}_0(\lambda) \quad \text{by Proposition 3.2.3} \\ &\cong \text{Par}(\lambda) \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda) \quad \text{by (3.2.3)} \\ &\cong \mathbb{B}^{\frac{\infty}{2}}(\lambda) \quad \text{by Proposition 3.2.4,} \end{aligned}$$

as desired.

In the remainder of this paper, we will give proofs of the results above; we prove Proposition 3.1.3 in §4.2, Theorem 3.1.5 in §4.3, Proposition 3.2.2 in §5, and Proposition 3.2.4 in §7.2.

4 Proofs of Proposition 3.1.3 and Theorem 3.1.5.

4.1 Some technical lemmas.

Lemma 4.1.1 ([BFP, Lemma 4.3]). *We have $\ell(r_\alpha) \leq 2\langle \alpha^\vee, \rho \rangle - 1$ for all $\alpha \in \Delta^+$.*

Lemma 4.1.2. *Let $x = vt_\zeta \in W_{\text{af}}$ with $v \in W$ and $\zeta \in Q^\vee$, and let $\beta = \alpha + n\delta \in \Delta_{\text{af}}^+$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}_{\geq 0}$. Then, $\ell^{\frac{\infty}{2}}(r_\beta x) > \ell^{\frac{\infty}{2}}(x)$ if and only if $v^{-1}\alpha$ is a positive root. In particular, if $\beta = \alpha_i$ for some $i \in I_{\text{af}}$ (note that $\alpha = \alpha_i$ and $n = 0$ if $i \in I$, and $\alpha = -\theta$ and $n = 1$ if $i = 0$), then*

$$\ell^{\frac{\infty}{2}}(r_i x) = \begin{cases} \ell^{\frac{\infty}{2}}(x) + 1 & \text{if } v^{-1}\alpha \text{ is a positive root,} \\ \ell^{\frac{\infty}{2}}(x) - 1 & \text{if } v^{-1}\alpha \text{ is a negative root.} \end{cases} \quad (4.1.1)$$

Proof. Assume that $\alpha \in \Delta$ is a negative root; note that $n \geq 1$. We see by (2.1.2) that $r_\beta x = r_\alpha t_{n\alpha^\vee} vt_\zeta = v r_{v^{-1}\alpha} t_{nv^{-1}\alpha^\vee + \zeta}$. If $v^{-1}\alpha$ is a positive root, then we have

$$\begin{aligned} \ell^{\frac{\infty}{2}}(r_\beta x) &= \ell(v r_{v^{-1}\alpha}) + 2\langle nv^{-1}\alpha^\vee + \zeta, \rho \rangle \geq \ell(v) - \ell(r_{v^{-1}\alpha}) + 2n\langle v^{-1}\alpha^\vee, \rho \rangle + 2\langle \zeta, \rho \rangle \\ &\geq \ell(v) - 2\langle v^{-1}\alpha^\vee, \rho \rangle + 1 + 2n\langle v^{-1}\alpha^\vee, \rho \rangle + 2\langle \zeta, \rho \rangle \quad \text{by Lemma 4.1.1} \\ &\geq \ell(v) + 2\langle \zeta, \rho \rangle + 1 \quad \text{since } v^{-1}\alpha \in \Delta^+ \text{ and } n \geq 1 \\ &= \ell^{\frac{\infty}{2}}(x) + 1 > \ell^{\frac{\infty}{2}}(x). \end{aligned}$$

If $v^{-1}\alpha$ is a negative root, then we have

$$\ell^{\frac{\infty}{2}}(r_\beta x) = \ell(v r_{v^{-1}\alpha}) + 2\langle nv^{-1}\alpha^\vee + \zeta, \rho \rangle \leq \ell(v) + \ell(r_{v^{-1}\alpha}) + 2n\langle v^{-1}\alpha^\vee, \rho \rangle + 2\langle \zeta, \rho \rangle$$

$$\begin{aligned}
&\leq \ell(v) + 2\langle v^{-1}\alpha^\vee, \rho \rangle - 1 + 2n\langle v^{-1}\alpha^\vee, \rho \rangle + 2\langle \zeta, \rho \rangle \quad \text{by Lemma 4.1.1} \\
&\leq \ell(v) + 2\langle \zeta, \rho \rangle - 1 \quad \text{since } -v^{-1}\alpha \in \Delta^+ \text{ and } n \geq 1 \\
&= \ell^{\frac{\infty}{2}}(x) - 1 < \ell^{\frac{\infty}{2}}(x).
\end{aligned}$$

The proof for the case that $\alpha \in \Delta$ is a positive root is easier.

In order to show (4.1.1), it suffices to verify that $\ell^{\frac{\infty}{2}}(r_i x) = \ell^{\frac{\infty}{2}}(x) \pm 1$. If $i \in I$, then it is obvious. Assume that $i = 0$. We see from [LNS³1, Proposition 5.11] that if $v^{-1}\theta$ is a negative root, then $\ell(r_\theta v) = \ell(v) - 2\langle -v^{-1}\theta^\vee, \rho \rangle + 1$. The desired equality above for $i = 0$ follows easily from this equality, together with the computation above. This proves the lemma. $\square$

Remark 4.1.3. Let $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. For each $x \in (W^J)_{\text{af}}$ and $i \in I_{\text{af}}$, we deduce from Lemma 2.3.6 that $r_i x \in (W^J)_{\text{af}}$ if and only if $\langle \alpha_i^\vee, x\lambda \rangle \neq 0$, and from (4.1.1) that

$$\begin{cases} x \xrightarrow{\alpha_i} r_i x & \text{in } \text{SiB}^J \iff \langle \alpha_i^\vee, x\lambda \rangle > 0 \iff x\lambda \xrightarrow{\alpha_i} r_i x\lambda & \text{in } \text{LZ}(\lambda), \\ r_i x \xrightarrow{\alpha_i} x & \text{in } \text{SiB}^J \iff \langle \alpha_i^\vee, x\lambda \rangle < 0 \iff r_i x\lambda \xrightarrow{\alpha_i} x\lambda & \text{in } \text{LZ}(\lambda). \end{cases} \quad (4.1.2)$$

We prove the “diamond lemma” for SiB^J (cf. [Li2, Lemma 4.1] and [LNS³1, Lemma 5.14]).

Lemma 4.1.4. *Let J be a subset of I . Let $x \in (W^J)_{\text{af}}$, $\beta \in \Delta_{\text{af}}^+$, and $i \in I_{\text{af}}$. Assume that $x \xrightarrow{\beta} r_\beta x =: y$ and $x \xrightarrow{\alpha_i} r_i x$ in SiB^J . If we write $y^{-1}\alpha_i \in \Delta_{\text{af}}$ as $y^{-1}\alpha_i = \gamma + n\delta$ with $\gamma \in \Delta$ and $n \in \mathbb{Z}$, then $\gamma \notin \Delta_J^+$. Moreover, if $\gamma \in \Delta^+ \setminus \Delta_J^+$, then $\beta \neq \alpha_i$, and we have dotted edges in SiB^J as in the following diagram:*

$$\begin{array}{ccc} x & \xrightarrow{\beta} & y \\ \alpha_i \downarrow & & \vdots \downarrow \alpha_i \\ r_i x & \xrightarrow{\dots \dots \dots r_i \beta} & r_i y \end{array}$$

If γ is a negative root, then $\beta = \alpha_i$.

Proof. First, suppose, for a contradiction, that $\gamma \in \Delta_J^+$. Then, we must have $y^{-1}\alpha_i \in \Delta_{\text{af}}^+$. Indeed, suppose that $y^{-1}\alpha_i$ is a negative real root. Since $\gamma \in \Delta_J^+$ by our assumption, we obtain $y^{-1}\alpha_i \in (\Delta_J)_{\text{af}} \cap (-\Delta_{\text{af}}^+)$. Since $y \in (W^J)_{\text{af}}$ by the assumption, it follows from the definition of $(W^J)_{\text{af}}$ that $\alpha_i = yy^{-1}\alpha_i$ is a negative real root, which is a contradiction. Thus, we have $y^{-1}\alpha_i \in \Delta_{\text{af}}^+$, and hence $y^{-1}\alpha_i \in (\Delta_J)_{\text{af}}^+$. Here, since $x \in (W^J)_{\text{af}}$, we see that $\Delta_{\text{af}}^+ \ni xy^{-1}\alpha_i = r_\beta \alpha_i = \alpha_i - \langle \beta^\vee, \alpha_i \rangle \beta$. Therefore, we deduce that $\langle \beta^\vee, \alpha_i \rangle \leq 0$; in particular, $\beta \neq \alpha_i$. We write $x^{-1}\alpha_i$ and $x^{-1}\beta$ as $x^{-1}\alpha_i = \gamma_1 + n_1\delta$ and $x^{-1}\beta = \gamma_2 + n_2\delta$, with $\gamma_1, \gamma_2 \in \Delta$ and $n_1, n_2 \in \mathbb{Z}$, respectively. Since we have $x \xrightarrow{\beta} r_\beta x = y$ and $x \xrightarrow{\alpha_i} r_i x$ in SiB^J by the assumption, we see by Lemma 4.1.2 that $\gamma_1, \gamma_2 \in \Delta^+$. Also, since $x, r_\beta x = xr_{x^{-1}\beta}$, and $r_i x = xr_{x^{-1}\alpha_i}$ are contained in

$(W^J)_{\text{af}}$, we see from Proposition 2.3.2 that $x^{-1}\beta, x^{-1}\alpha_i \notin (\Delta_J)_{\text{af}}$, and hence $\gamma_1, \gamma_2 \in \Delta^+ \setminus \Delta_J^+$. Therefore, we deduce that $\gamma_1 - \langle \beta^\vee, \alpha_i \rangle \gamma_2 \notin \Delta_J^+$, since $\langle \beta^\vee, \alpha_i \rangle \leq 0$ as seen above. Because

$$\begin{aligned} \gamma + n\delta &= y^{-1}\alpha_i = x^{-1}r_\beta\alpha_i = x^{-1}\alpha_i - \langle \beta^\vee, \alpha_i \rangle x^{-1}\beta \\ &= \{\gamma_1 - \langle \beta^\vee, \alpha_i \rangle \gamma_2\} + \{n_1 - \langle \beta^\vee, \alpha_i \rangle n_2\}\delta, \end{aligned}$$

we obtain $\gamma = \gamma_1 - \langle \beta^\vee, \alpha_i \rangle \gamma_2 \notin \Delta_J^+$, which contradicts our assumption that $\gamma \in \Delta_J^+$. Thus, we conclude that $\gamma \notin \Delta_J^+$, as desired.

Assume that $\gamma \in \Delta^+ \setminus \Delta_J^+$. Suppose, for a contradiction, that $\beta = \alpha_i$. Then we have $\gamma + n\delta = y^{-1}\alpha_i = x^{-1}r_\beta\alpha_i = x^{-1}r_i\alpha_i = -x^{-1}\alpha_i$. Since $x \xrightarrow{\alpha_i} r_ix$ in SiB^J by the assumption, it follows immediately from Lemma 4.1.2 that $x^{-1}\alpha_i = \gamma_1 + n_1\delta$ for some $\gamma_1 \in \Delta^+$ and $n_1 \in \mathbb{Z}$. Hence we obtain $\gamma = -\gamma_1 \in (-\Delta^+)$, which contradicts our assumption. Thus, we conclude that $\beta \neq \alpha_i$. Also, since $\langle \alpha_i^\vee, y\lambda \rangle = \langle y^{-1}\alpha_i^\vee, \lambda \rangle = \langle \gamma^\vee, \lambda \rangle > 0$, it follows immediately from (4.1.2) that $y \xrightarrow{\alpha_i} r_iy$ in SiB^J . Therefore, we have

$$\begin{aligned} \ell^{\frac{\infty}{2}}(r_iy) &= \underbrace{\ell^{\frac{\infty}{2}}(r_iy) - \ell^{\frac{\infty}{2}}(y)}_{=1} + \underbrace{\ell^{\frac{\infty}{2}}(y) - \ell^{\frac{\infty}{2}}(x)}_{=1} + \ell^{\frac{\infty}{2}}(x) \\ &= \ell^{\frac{\infty}{2}}(x) + 2 = \ell^{\frac{\infty}{2}}(r_ix) - 1 + 2 = \ell^{\frac{\infty}{2}}(r_ix) + 1. \end{aligned}$$

From this equation, we conclude that $r_ix \xrightarrow{r_i\beta} r_iy$.

Assume that γ is a negative root, and suppose, for a contradiction, that $\beta \neq \alpha_i$. Then we have $r_i\beta \in \Delta_{\text{af}}^+$. Since $x \xrightarrow{\beta} y$ in SiB^J by the assumption, it follows from Lemma 4.1.2 that $\ell^{\frac{\infty}{2}}(r_ix) < \ell^{\frac{\infty}{2}}(r_iy)$. Since $\langle \alpha_i^\vee, r_iy\lambda \rangle = -\langle y^{-1}\alpha_i^\vee, \lambda \rangle = -\langle \gamma^\vee, \lambda \rangle > 0$, we see by (4.1.2) that $r_iy \xrightarrow{\alpha_i} y$ in SiB^J . Therefore, we have $\ell^{\frac{\infty}{2}}(x) < \ell^{\frac{\infty}{2}}(r_ix) < \ell^{\frac{\infty}{2}}(r_iy) < \ell^{\frac{\infty}{2}}(y)$, which contradicts the equality $\ell^{\frac{\infty}{2}}(y) = \ell^{\frac{\infty}{2}}(x) + 1$. This completes the proof of the lemma. $\square$

The following lemma can be proved in exactly the same way as Lemma 4.1.4.

Lemma 4.1.5. *Let J be a subset of I . Let $x \in (W^J)_{\text{af}}$, $\beta \in \Delta_{\text{af}}^+$, and $i \in I$. Assume that $y := r_\beta x \xrightarrow{\beta} x$ and $r_ix \xrightarrow{\alpha_i} x$ in SiB^J . If we write $y^{-1}\alpha_i \in \Delta_{\text{af}}$ as $y^{-1}\alpha_i = -\gamma + n\delta$ with $\gamma \in \Delta$ and $n \in \mathbb{Z}$, then $\gamma \notin \Delta_J^+$. Moreover, if $\gamma \in \Delta^+ \setminus \Delta_J^+$, then $\beta \neq \alpha_i$, and we have dotted edges in SiB^J as in the following diagram:*

$$\begin{array}{ccc} r_iy & \cdots \xrightarrow{\beta} & r_ix \\ \vdots & & \downarrow \alpha_i \\ y & \xrightarrow{\beta} & x \end{array}$$

If γ is a negative root, then $\beta = \alpha_i$.

An inductive argument, which uses Lemmas 4.1.4 and 4.1.5, proves the following proposition.

Lemma 4.1.6. Let $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. Let $0 < a \leq 1$ be a rational number. Let $x, y \in (W^J)_{\text{af}}$, and assume that there exists a directed path $x = y_0 \xrightarrow{\beta_1} y_1 \xrightarrow{\beta_2} \cdots \xrightarrow{\beta_k} y_k = y$ from x to y in $\text{SiB}(\lambda; a)$. Let $i \in I_{\text{af}}$.

- (1) If $\langle \alpha_i^\vee, y_m \lambda \rangle > 0$ for all $0 \leq m \leq k$, or if $\langle \alpha_i^\vee, y_m \lambda \rangle < 0$ for all $0 \leq m \leq k$, then there exists a directed path from $r_i x$ to $r_i y$ in $\text{SiB}(\lambda; a)$ of the form:

$$r_i x = r_i y_0 \xrightarrow{r_i \beta_1} r_i y_1 \xrightarrow{r_i \beta_2} \cdots \xrightarrow{r_i \beta_k} r_i y_k = r_i y.$$

- (2) Assume that $\langle \alpha_i^\vee, x \lambda \rangle > 0$, and $\langle \alpha_i^\vee, y_m \lambda \rangle \leq 0$ for some $1 \leq m \leq k$; let l denote the minimum of all such m 's. Then, $\beta_l = \alpha_i$, and there exists a directed path from $r_i x$ to y in $\text{SiB}(\lambda; a)$ of the form:

$$r_i x = r_i y_0 \xrightarrow{r_i \beta_1} \cdots \xrightarrow{r_i \beta_{l-1}} r_i y_{l-1} = y_l \xrightarrow{\beta_{l+1}} \cdots \xrightarrow{\beta_k} y_k = y.$$

- (3) Assume that $\langle \alpha_i^\vee, y \lambda \rangle < 0$, and $\langle \alpha_i^\vee, y_m \lambda \rangle \geq 0$ for some $0 \leq m \leq k-1$; let l denote the maximum of all such m 's. Then, $\beta_{l+1} = \alpha_i$, and there exists a directed path from x to $r_i y$ in $\text{SiB}(\lambda; a)$ of the form:

$$x = y_0 \xrightarrow{\beta_1} \cdots \xrightarrow{\beta_l} y_l = r_i y_{l+1} \xrightarrow{r_i \beta_{l+2}} \cdots \xrightarrow{r_i \beta_k} r_i y_k = r_i y.$$

Lemma 4.1.7. Let J be a subset of I . Let $\xi \in Q^{\vee, J\text{-ad}}$, and $\beta \in \Delta_{\text{af}}^+$. If $z_\xi t_\xi \xrightarrow{\beta} r_\beta z_\xi t_\xi$ in SiB^J , then $\beta = \alpha_i$ for some $i \in I \setminus J$.

Proof. Write β as $\beta = \alpha + n\delta$, with $\alpha \in \Delta$ and $n \in \mathbb{Z}_{\geq 0}$. Since $z_\xi t_\xi, r_\beta z_\xi t_\xi \in (W^J)_{\text{af}}$ by the assumption, we see by Lemma 2.3.7 that $r_\beta \in (W^J)_{\text{af}}$, which implies that $\beta \notin (\Delta_J)_{\text{af}}$ and hence $\alpha \notin \Delta_J$. Also, since $\ell^{\frac{\infty}{2}}(r_\beta z_\xi t_\xi) = \ell^{\frac{\infty}{2}}(z_\xi t_\xi) + 1 > \ell^{\frac{\infty}{2}}(z_\xi t_\xi)$ by the assumption, we see from Lemma 4.1.2 that $z_\xi^{-1} \alpha$ is a positive root. Because $\alpha \notin \Delta_J$ as seen above, and $z_\xi \in W_J$, we deduce that $\alpha \in \Delta^+$, and hence $\alpha \in \Delta^+ \setminus \Delta_J^+$. We claim that $r_\alpha \in W^J$. Indeed, if $n = 0$, then we have $r_\alpha = r_\beta \in (W^J)_{\text{af}}$. Hence we obtain $r_\alpha = \Pi^J(r_\alpha) = \lfloor r_\alpha \rfloor$ by Lemma 2.3.3 (1), which implies that $r_\alpha \in W^J$. Assume that $n \geq 1$. By (2.1.2), we have $r_\beta = r_\alpha t_{n\alpha^\vee}$. Since $r_\beta \in (W^J)_{\text{af}}$ as seen above, we have $n\alpha^\vee \in Q^{\vee, J\text{-ad}}$ by (2.3.5). Therefore, we have $\langle n\alpha^\vee, \gamma \rangle \in \{0, -1\}$ for all $\gamma \in \Delta_J^+$, and in particular, $\langle \alpha^\vee, \gamma \rangle \leq 0$ for all $\gamma \in \Delta_J^+$. From this, we see that $r_\alpha \gamma = \gamma - \langle \alpha^\vee, \gamma \rangle \alpha \in \Delta^+$ for all $\gamma \in \Delta_J^+$ since $\alpha \in \Delta^+$, which implies that $r_\alpha \in W^J$ by (2.1.3), as claimed. Now we compute

$$\begin{aligned} \ell^{\frac{\infty}{2}}(r_\beta z_\xi t_\xi) &= \ell^{\frac{\infty}{2}}(r_\alpha z_\xi t_{\xi + n z_\xi^{-1} \alpha^\vee}) = \ell(r_\alpha z_\xi) + 2\langle \xi + n z_\xi^{-1} \alpha^\vee, \rho \rangle \\ &= \ell(r_\alpha) + \underbrace{\ell(z_\xi) + 2\langle \xi, \rho \rangle}_{=\ell^{\frac{\infty}{2}}(z_\xi t_\xi)} + 2n\langle z_\xi^{-1} \alpha^\vee, \rho \rangle \quad \text{since } r_\alpha \in W^J \text{ and } z_\xi \in W_J. \end{aligned}$$

Since $\ell^{\frac{\infty}{2}}(r_\beta z_\xi t_\xi) = \ell^{\frac{\infty}{2}}(z_\xi t_\xi) + 1$ by the assumption, it follows from this equation that $\ell(r_\alpha) + 2n\langle z_\xi^{-1} \alpha^\vee, \rho \rangle = 1$. Here, recall that $z_\xi^{-1} \alpha$ is a positive root, and that $n \geq 0$. Therefore, we obtain $\ell(r_\alpha) = 1$ and $n = 0$, which implies that $\beta = \alpha_i$ for some $i \in I \setminus J$. This proves the lemma. $\square$

4.2 Proof of Proposition 3.1.3.

In this subsection, we fix $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$.

Let $\eta = (x_1, x_2, \dots, x_s; a_0, a_1, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$; recall that

$$\bar{\eta} = (x_1\lambda, x_2\lambda, \dots, x_s\lambda; a_0, a_1, \dots, a_s).$$

Hence it suffices to show that $x_u\lambda > x_{u+1}\lambda$, and that there exists a directed path from $x_{u+1}\lambda$ to $x_u\lambda$ in $\text{LZ}(\lambda; a_u)$. This follows immediately from the next proposition.

Proposition 4.2.1. *Let $0 < a \leq 1$ be a rational number, $x \in (W^J)_{\text{af}}$, and $\beta \in \Delta_{\text{af}}^+$. Then, $x \xrightarrow{\beta} r_\beta x$ in $\text{SiB}(\lambda; a)$ if and only if $x\lambda \xrightarrow{\beta} r_\beta x\lambda$ in $\text{LZ}(\lambda; a)$.*

Proof. In view of conditions (2.2.1) and (3.1.1), we need only show that

$$x \xrightarrow{\beta} r_\beta x \text{ in } \text{SiB}^J \text{ if and only if } x\lambda \xrightarrow{\beta} r_\beta x\lambda \text{ in } \text{LZ}(\lambda); \quad (4.2.1)$$

write $\beta \in \Delta_{\text{af}}^+$ as $\beta = \alpha + n\delta$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}_{\geq 0}$. We deduce from [AK, Lemma 1.4] that there exist $i_1, i_2, \dots, i_k \in I_{\text{af}}$ such that

$$x\lambda = \mu_k \xrightarrow{\alpha_{i_k}} \mu_{k-1} \xrightarrow{\alpha_{i_{k-1}}} \dots \xrightarrow{\alpha_{i_2}} \mu_1 \xrightarrow{\alpha_{i_1}} \mu_0 \quad \text{in } \text{LZ}(\lambda), \quad (4.2.2)$$

with $\mu_0 \equiv \lambda \pmod{\mathbb{Q}\delta}$, where $\mu_m = r_{i_{m+1}} \cdots r_{i_k} x\lambda$ for $0 \leq m \leq k$. We will show (4.2.1) by induction on the length k of the directed path (4.2.2).

Step 1. If $k = 0$, then we have $x\lambda \equiv \lambda \pmod{\mathbb{Q}\delta}$, which implies that $x \in (W^J)_{\text{af}}$ is of the form $x = z_\xi t_\xi$ for some $\xi \in Q^\vee, J\text{-ad}$ (see (2.3.5)). First, let us show the “only if” part of (4.2.1) in this case. By Lemma 4.1.7, we have $\beta = \alpha_i$ for some $i \in I \setminus J$. Hence it follows immediately from (4.1.2) that $x\lambda \xrightarrow{\alpha_i = \beta} r_i x\lambda = r_\beta x\lambda$ in $\text{LZ}(\lambda)$.

Next, let us show the “if part” of (4.2.1) in the case that $x = z_\xi t_\xi$ with $\xi \in Q^\vee, J\text{-ad}$. We see from [NS4, Lemma 2.11] that β is of the form $\beta = \alpha$ with $\alpha \in \Delta^+$, or $\beta = -\alpha + \delta$ with $\alpha \in \Delta^+$. Since $\langle \beta^\vee, x\lambda \rangle > 0$, and $x\lambda \equiv \lambda \pmod{\mathbb{Q}\delta}$, it follows immediately that $\beta = \alpha \in \Delta^+ \setminus \Delta_J^+$. Let $[r_\alpha] = r_{j_p} \cdots r_{j_1}$ be a reduced expression of $[r_\alpha] \in W$; since $x\lambda \equiv \lambda \pmod{\mathbb{Q}\delta}$, we have

$$\langle \alpha_{j_q}^\vee, r_{j_{q-1}} \cdots r_{j_1} x\lambda \rangle = \langle \alpha_{j_q}^\vee, r_{j_{q-1}} \cdots r_{j_1} \lambda \rangle > 0$$

for all $1 \leq q \leq p$. Then, by (4.1.2), we deduce that

$$x\lambda \xrightarrow{\alpha_{j_1}} r_{j_1} x\lambda \xrightarrow{\alpha_{j_2}} \dots \xrightarrow{\alpha_{j_p}} r_{j_p} \cdots r_{j_1} x\lambda = [r_\alpha] x\lambda \quad \text{in } \text{LZ}(\lambda). \quad (4.2.3)$$

Noting that $x\lambda \equiv \lambda \pmod{\mathbb{Q}\delta}$, we see that $[r_\alpha] x\lambda = r_\alpha x\lambda = r_\beta x\lambda$, and hence (4.2.3) is a directed path from $x\lambda$ to $r_\beta x\lambda$. However, we have $x\lambda \xrightarrow{\beta} r_\beta x\lambda$ by the assumption. From these, we deduce that $p = 1$, and hence $[r_\alpha] = r_i$ for some $i \in I$. Since $\langle \alpha^\vee, \lambda \rangle = \langle \beta^\vee, x\lambda \rangle > 0$ and $r_\alpha \lambda = r_i \lambda$, we see that $\beta = \alpha = \alpha_i$ (see Remark 2.2.2). Therefore, it follows from (4.1.2) that $x \xrightarrow{\alpha_i = \beta} r_i x = r_\beta x$ in SiB^J .

Step 2. Assume that the length k of the directed path (4.2.2) is greater than 0; for simplicity of notation, we set $i := i_k$ in (4.2.2). Since we have $x\lambda \xrightarrow{\alpha_i} r_i x\lambda$ in $\text{LZ}(\lambda)$ as the initial edge of (4.2.2), we have $\langle \alpha_i^\vee, x\lambda \rangle > 0$. Hence it follows from (4.1.2) that $x \xrightarrow{\alpha_i} r_i x$ in SiB^J .

First, let us show the “only if” part of (4.2.1). Set $y := r_\beta x$, and write $y^{-1}\alpha_i$ as $y^{-1}\alpha_i = \gamma + m\delta$, with $\gamma \in \Delta$ and $m \in \mathbb{Z}$; we see by Lemma 4.1.4 that $\gamma \notin \Delta_J^+$. If γ is a negative root, then $\beta = \alpha_i$ by Lemma 4.1.4, and hence we have $x\lambda \xrightarrow{\alpha_i=\beta} r_i x\lambda = r_\beta x\lambda$. Assume that $\gamma \in \Delta^+ \setminus \Delta_J^+$. Then, by Lemma 4.1.4, $\beta \neq \alpha_i$, and we obtain the following diagram in SiB^J :

$$\begin{array}{ccc} x & \xrightarrow{\beta} & y \\ \alpha_i \downarrow & & \vdots \downarrow \alpha_i \\ r_i x & \xrightarrow{\dots \dots r_i \beta} & r_i y \end{array}$$

By our induction hypothesis applied to $r_i x$, we have $r_i x\lambda \xrightarrow{r_i \beta} r_i y\lambda$ in $\text{LZ}(\lambda)$. Also, since $y \xrightarrow{\alpha_i} r_i y$ in SiB^J , it follows from (4.1.2) that $y\lambda \xrightarrow{\alpha_i} r_i y\lambda$ in $\text{LZ}(\lambda)$. Because $x\lambda \xrightarrow{\alpha_i} r_i x\lambda$ in $\text{LZ}(\lambda)$ as the initial edge of (4.2.2), we deduce that $x\lambda \xrightarrow{\beta} y\lambda$ by [Li2, Lemma 4.1 c)], as desired.

Next, let us show the “if” part of (4.2.1). Recall that $\langle \alpha_i^\vee, x\lambda \rangle > 0$. If $\langle \alpha_i^\vee, y\lambda \rangle \leq 0$, then it follows from [Li2, Corollary 1 in §4] that $\beta = \alpha_i$, and hence $x \xrightarrow{\alpha_i=\beta} r_i x = r_\beta x$. Assume that $\langle \alpha_i^\vee, y\lambda \rangle > 0$. Then we have $y \xrightarrow{\alpha_i} r_i y$ by (4.1.2). It follows immediately from [Li2, Lemma 4.1 c)] that $\beta \neq \alpha_i$, and $r_i x\lambda \xrightarrow{r_i \beta} r_i y\lambda$. By our induction hypothesis applied to $r_i x$, we have $r_i x \xrightarrow{r_i \beta} r_i y$. Therefore, we deduce from Lemma 4.1.5 that $x \xrightarrow{\beta} y$, as desired. This completes the proof of the proposition. $\square$

The next corollary follows immediately from Proposition 4.2.1 and [NS4, Lemma 2.11].

Corollary 4.2.2. *Let J be a subset of I . Let $x \in (W^J)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$ be such that $x \xrightarrow{\beta} r_\beta x$. Then, β is either of the following forms: $\beta = \alpha$ with $\alpha \in \Delta^+$, or $\beta = \alpha + \delta$ with $-\alpha \in \Delta^+$. Moreover, if $x = wz_\xi t_\xi$ with $w \in W^J$ and $\xi \in Q^\vee, J\text{-ad}$ (see (2.3.5)), then $w^{-1}\alpha \in \Delta^+ \setminus \Delta_J^+$ in both cases.*

4.3 Proof of Theorem 3.1.5.

We prove part (1) only for e_i ; the proof for f_i is similar. Let $\eta = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$ be such that $m_i^{\bar{\eta}} \leq -1$. Define $t_0, t_1 \in [0, 1]$ by (2.2.4) (for $\bar{\eta}$ and $i \in I_{\text{af}}$), and let $1 \leq p \leq q \leq s$ be such that $a_{p-1} \leq t_0 < a_q$ and $t_1 = a_q$. By the definition of e_i , we have

$$0 = a_0 \xrightarrow{x_1} \dots \xrightarrow{x_{p-1}} a_{p-1} \xrightarrow{x_p} \underbrace{t_0 \xrightarrow{r_i x_p} a_p \xrightarrow{r_i x_{p+1}} \dots \xrightarrow{r_i x_q} a_q}_{\text{“reflected” by } r_i} = t_1 \xrightarrow{x_{q+1}} \dots \xrightarrow{x_s} a_s = 1;$$

if $t_0 = a_{p-1}$, then we drop x_p and a_{p-1} , and if $r_i x_q = x_{q+1}$, then we drop x_{q+1} and $a_q = t_1$. We need to show that

- (i) $r_i x_u \in (W^J)_{\text{af}}$ for all $p \leq u \leq q$;
- (ii) if $t_0 \neq a_{p-1}$ (resp., $t_0 = a_{p-1}$ and $p > 1$), then there exists a directed path from $r_i x_p$ to x_p (resp., to x_{p-1}) in $\text{SiB}(\lambda; t_0)$;
- (iii) for each $p \leq u \leq q-1$, there exists a directed path from $r_i x_{u+1}$ to $r_i x_u$ in $\text{SiB}(\lambda; a_u)$;
- (iv) if $r_i x_q \neq x_{q+1}$, then there exists a directed path from x_{q+1} to $r_i x_q$ in $\text{SiB}(\lambda; t_1) = \text{SiB}(\lambda; a_q)$.

Proof of (i). As mentioned in Definition 2.2.6 (1), the function $H_i^{\bar{\eta}}(t)$ is strictly decreasing on $[t_0, t_1]$. Therefore, we see that $\langle \alpha_i^\vee, x_u \lambda \rangle < 0$ for all $p \leq u \leq q$. This implies that $x_u^{-1} \alpha_i \notin (\Delta_J)_{\text{af}}$, and hence $r_i x_u \in (W^J)_{\text{af}}$ by Lemma 2.3.6. ■

Proof of (ii). Since $\langle \alpha_i^\vee, x_p \lambda \rangle < 0$ as above, we have $r_i x_p \xrightarrow{\alpha_i} x_p$ in SiB^J by (4.1.2). By applying [Li2, Lemma 4.5 c)] to $\bar{\eta} \in \mathbb{B}(\lambda)$ and $t_0 \in [0, 1]$, we deduce that $t_0 \langle \alpha_i^\vee, x_p \lambda \rangle \in \mathbb{Z}$, which implies that the edge $r_i x_p \xrightarrow{\alpha_i} x_p$ above is an edge in $\text{LZ}(\lambda; t_0)$. Thus we have shown (ii) in the case that $t_0 \neq a_{p-1}$. Assume next that $t_0 = a_{p-1}$ and $p > 1$. By the assumption, there exists a directed path from x_p to x_{p-1} in $\text{SiB}(\lambda; a_{p-1}) = \text{SiB}(\lambda; t_0)$. By concatenating this directed path with $r_i x_p \xrightarrow{\alpha_i} x_p$ obtained above, we obtain a directed path from $r_i x_p$ to x_{p-1} in $\text{SiB}(\lambda; t_0)$. Thus we have shown (ii). ■

Proof of (iii). Fix $p \leq u \leq q-1$. Let $x_{u+1} = y_0 \xrightarrow{\beta_1} y_1 \xrightarrow{\beta_2} \cdots \xrightarrow{\beta_k} y_k = x_u$ be a directed path from x_{u+1} to x_u in $\text{SiB}(\lambda; a_u)$. Since $H_i^{\bar{\eta}}(t)$ is strictly decreasing on $[t_0, t_1]$, we see that $\langle \alpha_i^\vee, x_{u+1} \lambda \rangle < 0$. Also, since $H_i^{\bar{\eta}}(t_0) = m_i^{\bar{\eta}} + 1$ and $H_i^{\bar{\eta}}(t_1) = m_i^{\bar{\eta}}$, we see that $H_i^{\bar{\eta}}(a_u) \notin \mathbb{Z}$. Therefore, it follows from [Li2, Remark 4.6] that $\langle \alpha_i^\vee, y_l \lambda \rangle > 0$ for all $0 \leq l \leq k$, and hence that there exists a directed path from $r_i x_{u+1}$ to $r_i x_u$ in $\text{SiB}(\lambda; a_u)$ by Lemma 4.1.6 (1). ■

Proof of (iv). By the definition, there exists a directed path from x_{q+1} to x_q in $\text{SiB}(\lambda; a_q)$. By the definition of t_1 , we see that $\langle \alpha_i^\vee, x_{q+1} \lambda \rangle \geq 0$ and $\langle \alpha_i^\vee, x_q \lambda \rangle < 0$. Therefore, it follows from Lemma 4.1.6 (3) that there exists a directed path from x_{q+1} to $r_i x_q$ in $\text{SiB}(\lambda; a_q)$. ■

Thus we have proved part (1) of Theorem 3.1.5. Next, let us prove part (2). We see from the definition of root operators that for all $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$,

$$\begin{cases} \text{wt}(\eta) = \text{wt}(\bar{\eta}), \\ \overline{e_i \eta} = e_i \bar{\eta}, \overline{f_i \eta} = f_i \bar{\eta}, \\ \varepsilon_i(\eta) = \varepsilon_i(\bar{\eta}), \varphi_i(\eta) = \varphi_i(\bar{\eta}), \end{cases} \quad (4.3.1)$$

where we understand that $\bar{\mathbf{0}} = \mathbf{0}$. From (4.3.1), we can easily deduce that the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ satisfies the axioms for crystals, except for the axiom that

$$e_i \eta_1 = \eta_2 \text{ if and only if } \eta_1 = f_i \eta_2 \text{ for } \eta_1, \eta_2 \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \text{ and } i \in I_{\text{af}}. \quad (4.3.2)$$

Now we define $t_0^1, t_1^1 \in [0, 1]$ (resp., $t_0^2, t_1^2 \in [0, 1]$) by (2.2.4) (resp., (2.2.5)) for $\overline{\eta}_1$ (resp., $\overline{\eta}_2$) and $i \in I_{\text{af}}$; note that $H_i^{\overline{\eta}_1}(t)$ (resp., $H_i^{\overline{\eta}_2}(t)$) is strictly decreasing (resp., increasing) on $[t_0^1, t_1^1]$ (resp., $[t_0^2, t_1^2]$). Then we deduce from the definitions that $t_0^1 = t_0^2$ and $t_1^1 = t_1^2$. Therefore, (4.3.2) follows immediately from the definition of the root operators e_i and f_i . This completes the proof of Theorem 3.1.5.

Remark 4.3.1.

- (1) By (4.3.1), the map $\bar{\cdot} : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \rightarrow \mathbb{B}(\lambda)$, $\eta \mapsto \overline{\eta}$, is a strict morphism of crystals in the sense of [Kas1, §1.5]; also, this map is surjective.
- (2) By (4.3.1) and [Li2, Lemma 2.1 c)], we have $\varepsilon_i(\eta) = -m_i^{\overline{\eta}}$ and $\varphi_i(\eta) = H_i^{\overline{\eta}}(1) - m_i^{\overline{\eta}}$ for all $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$.

5 Proof of Proposition 3.2.2.

Throughout this section, we fix $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$.

5.1 Extremal elements in $\mathcal{B}_0(\lambda)$ and $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$.

We know from [Kas1, §7] that the affine Weyl group W_{af} acts on $\mathcal{B}(\lambda)$ by

$$r_i b := \begin{cases} f_i^n b & \text{if } n = \langle \alpha_i^\vee, \text{wt}(b) \rangle \geq 0, \\ e_i^{-n} b & \text{if } n = \langle \alpha_i^\vee, \text{wt}(b) \rangle \leq 0, \end{cases} \quad (5.1.1)$$

for each $b \in \mathcal{B}(\lambda)$ and $i \in I_{\text{af}}$.

Proposition 5.1.1 (cf. [Kas5, Conjecture 5.11]; see also Remark 2.3.1). *The following equality holds:*

$$(W_J)_{\text{af}} = \{x \in W_{\text{af}} \mid xu_\lambda = u_\lambda\}.$$

Proof. Write $\lambda \in P^+$ as $\lambda = \sum_{i \in I} m_i \varpi_i$ with $m_i \in \mathbb{Z}_{\geq 0}$, $i \in I$. Then we know from [BN, Remark 4.17] (see also [Kas5, §13]) that there exists an embedding $\Psi : \mathcal{B}_0(\lambda) \hookrightarrow \bigotimes_{i \in I} \mathcal{B}(\varpi_i)^{\otimes m_i}$ of crystals such that $\Psi(u_\lambda) = \bigotimes_{i \in I} u_{\varpi_i}^{\otimes m_i}$. Recall that for each $i \in I$, $u_{\varpi_i} \in \mathcal{B}(\varpi_i)$ is an extremal element of weight ϖ_i in the sense of [Kas1, Definition 8.1.1]; in particular, we have $\text{wt}(xu_{\varpi_i}) = x\varpi_i$, $\varepsilon_j(xu_{\varpi_i}) = \max\{0, -\langle \alpha_j^\vee, x\varpi_i \rangle\}$, and $\varphi_j(xu_{\varpi_i}) = \max\{0, \langle \alpha_j^\vee, x\varpi_i \rangle\}$ for every $x \in W_{\text{af}}$ and $j \in I_{\text{af}}$. From these, using the tensor product rule for crystals, we can show by induction on $\ell(x)$ (see also [AK, Lemma 1.6]) that $\Psi(xu_\lambda) = \bigotimes_{i \in I} (xu_{\varpi_i})^{\otimes m_i}$ for all $x \in W_{\text{af}}$. Therefore, we deduce that

$$\{x \in W_{\text{af}} \mid xu_\lambda = u_\lambda\} = \bigcap_{i \in I \setminus J} \{x \in W_{\text{af}} \mid xu_{\varpi_i} = u_{\varpi_i}\}. \quad (5.1.2)$$

Also, we know from [Kas5, Lemma 5.6] that

$$\{x \in W_{\text{af}} \mid xu_{\varpi_i} = u_{\varpi_i}\} = \langle r_\beta \mid \beta \in \Delta_{\text{af}}^+, \langle \beta^\vee, \varpi_i \rangle = 0 \rangle. \quad (5.1.3)$$

If $\beta \in (\Delta_J)_{\text{af}}^+$, then $\langle \beta^\vee, \varpi_i \rangle = 0$, and hence $r_\beta \in \{x \in W_{\text{af}} \mid xu_{\varpi_i} = u_{\varpi_i}\}$ for all $i \in I \setminus J$. Because $(W_J)_{\text{af}} = \langle r_\beta \mid \beta \in (\Delta_J)_{\text{af}}^+ \rangle$ by Remark 2.3.1, it follows immediately that $(W_J)_{\text{af}} \subset \{x \in W_{\text{af}} \mid xu_\lambda = u_\lambda\}$.

Let us show the opposite inclusion. Let $x \in W_{\text{af}}$ be such that $xu_\lambda = u_\lambda$. By (5.1.2), we have $xu_{\varpi_i} = u_{\varpi_i}$ for all $i \in I \setminus J$; in particular, $x\varpi_i = \varpi_i$ for all $i \in I \setminus J$ since the weight of xu_{ϖ_i} is equal to $x\varpi_i$. We write x as $x = wt_\xi$ with $w \in W$ and $\xi \in Q^\vee$. Then, for all $i \in I \setminus J$, we have $\varpi_i = x\varpi_i = w\varpi_i - \langle \xi, \varpi_i \rangle \delta$ by (2.1.1), and hence $w\varpi_i = \varpi_i$ and $\langle \xi, \varpi_i \rangle = 0$. Therefore, we deduce that $w \in W_J \subset (W_J)_{\text{af}}$, and $\xi \in Q_J^\vee$, which implies that $t_\xi \in (W_J)_{\text{af}}$. Thus we obtain $x \in (W_J)_{\text{af}}$, and hence $(W_J)_{\text{af}} \supset \{x \in W_{\text{af}} \mid xu_\lambda = u_\lambda\}$. This completes the proof of the proposition. $\square$

Recall that $u_\lambda \in \mathcal{B}_0(\lambda)$ is an extremal element of weight λ in the sense of [Kas1, Definition 8.1.1]. From Proposition 5.1.1, we see that the set $\{yu_\lambda \mid y \in W_{\text{af}}\}$ is in bijective correspondence with the quotient set $W_{\text{af}}/(W_J)_{\text{af}}$, and hence with the set $(W^J)_{\text{af}}$ by Proposition 2.3.2. Hence we set $u_x := xu_\lambda$ for $x \in (W^J)_{\text{af}}$; remark that

$$\text{wt}(u_x) = x\lambda, \quad \varepsilon_i(u_x) = \max\{0, -\langle \alpha_i^\vee, x\lambda \rangle\}, \quad \varphi_i(u_x) = \max\{0, \langle \alpha_i^\vee, x\lambda \rangle\} \quad (5.1.4)$$

for all $x \in (W^J)_{\text{af}}$ and $i \in I_{\text{af}}$. It follows from Proposition 5.1.1 that $yu_\lambda = u_{\Pi^J(y)}$ for all $y \in W_{\text{af}}$.

Now, we set $\eta_x := (x; 0, 1) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ for $x \in (W^J)_{\text{af}}$. By Remark 4.3.1 (2) and (5.1.4), we see that for each $x \in (W^J)_{\text{af}}$ and $i \in I_{\text{af}}$,

$$\begin{aligned} \text{wt}(\eta_x) &= x\lambda = \text{wt}(u_x), & \varepsilon_i(\eta_x) &= \max\{0, -\langle \alpha_i^\vee, x\lambda \rangle\} = \varepsilon_i(u_x), \\ \varphi_i(\eta_x) &= \max\{0, \langle \alpha_i^\vee, x\lambda \rangle\} = \varphi_i(u_x). \end{aligned} \quad (5.1.5)$$

Next, for $x \in (W^J)_{\text{af}}$ and $i \in I_{\text{af}}$, we set

$$r_i \eta_x := \begin{cases} f_i^n \eta_x & \text{if } n = \langle \alpha_i^\vee, x\lambda \rangle \geq 0, \\ e_i^{-n} \eta_x & \text{if } n = \langle \alpha_i^\vee, x\lambda \rangle \leq 0; \end{cases}$$

observe that $r_i \eta_x \neq \mathbf{0}$ for any $x \in W_{\text{af}}$ and $i \in I_{\text{af}}$. Then we have

$$r_{i_p} r_{i_{p-1}} \cdots r_{i_2} r_{i_1} \eta_x = \eta_{\Pi^J(r_{i_p} r_{i_{p-1}} \cdots r_{i_2} r_{i_1} x)} \quad (5.1.6)$$

for all $x \in W_{\text{af}}$ and $i_1, \dots, i_p \in I_{\text{af}}$. Let us show (5.1.6) by induction on p . If $p = 0$, then (5.1.6) is obvious. Assume that $p > 0$. By our induction hypothesis, we have $r_{i_{p-1}} \cdots r_{i_2} r_{i_1} \eta_x = \eta_{\Pi^J(r_{i_{p-1}} \cdots r_{i_2} r_{i_1} x)}$; for simplicity of notation, we set $y := \Pi^J(r_{i_{p-1}} \cdots r_{i_2} r_{i_1} x) \in (W^J)_{\text{af}}$. If $n =$

$\langle \alpha_{i_p}^\vee, y\lambda \rangle > 0$, then $y^{-1}\alpha_{i_p} \notin (\Delta_J)_{\text{af}}$, and hence $r_{i_p}y \in (W^J)_{\text{af}}$ by Lemma 2.3.6. Therefore, we obtain

$$r_{i_p}y = \Pi^J(r_{i_p}y) = \Pi^J(r_{i_p}\Pi^J(r_{i_{p-1}} \cdots r_{i_2}r_{i_1}x)) = \Pi^J(r_{i_p}r_{i_{p-1}} \cdots r_{i_2}r_{i_1}x).$$

Also, it is easily shown by induction on k that $f_{i_p}^k \eta_y = (r_{i_p}y, y; 0, k/n, 1)$ for $0 \leq k \leq n$; in particular, we obtain $f_{i_p}^n \eta_y = \eta_{r_{i_p}y}$. From these, we deduce that

$$r_{i_p}r_{i_{p-1}} \cdots r_{i_2}r_{i_1}\eta_x = r_{i_p}\eta_y = f_{i_p}^n \eta_y = \eta_{r_{i_p}y} = \eta_{\Pi^J(r_{i_p}r_{i_{p-1}} \cdots r_{i_2}r_{i_1}x)},$$

as desired. The proof for the case that $n = \langle \alpha_{i_p}^\vee, y\lambda \rangle < 0$ is similar. If $n = \langle \alpha_{i_p}^\vee, y\lambda \rangle = 0$, then we have $r_{i_p}\eta_y = \eta_y$ by the definition. Also, we see that $y^{-1}\alpha_{i_p} \in (\Delta_J)_{\text{af}}$, and hence $r_{y^{-1}\alpha_{i_p}} \in (W_J)_{\text{af}}$ (see Remark 2.3.1). Hence it follows that $\Pi^J(r_{i_p}y) = \Pi^J(yr_{y^{-1}\alpha_{i_p}}) = \Pi^J(y) = y$. Therefore, we deduce that

$$r_{i_p}r_{i_{p-1}} \cdots r_{i_2}r_{i_1}\eta_x = r_{i_p}\eta_y = \eta_y = \eta_{\Pi^J(r_{i_p}y)} = \eta_{\Pi^J(r_{i_p}r_{i_{p-1}} \cdots r_{i_2}r_{i_1}x)},$$

as desired.

Remark 5.1.2. It follows from (5.1.6) that $\eta_x \in \mathbb{B}_0^{\infty}(\lambda)$ for all $x \in (W^J)_{\text{af}}$.

5.2 N -multiple maps.

Proposition 5.2.1. *Let $N \in \mathbb{Z}_{>0}$. There exists a unique injective map $\sigma_N : \mathcal{B}_0(\lambda) \hookrightarrow \mathcal{B}_0(\lambda)^{\otimes N}$ such that $\sigma_N(u_\lambda) = u_\lambda^{\otimes N}$, and*

$$\text{wt}(\sigma_N(b)) = N \text{wt}(b), \quad \varepsilon_i(\sigma_N(b)) = N \varepsilon_i(b), \quad \varphi_i(\sigma_N(b)) = N \varphi_i(b), \quad (5.2.1)$$

$$\sigma_N(e_i b) = e_i^N \sigma_N(b), \quad \sigma_N(f_i b) = f_i^N \sigma_N(b), \quad (5.2.2)$$

for $b \in \mathcal{B}_0(\lambda)$ and $i \in I_{\text{af}}$, where we understand that $\sigma_N(\mathbf{0}) = \mathbf{0}$.

Proof. We know from [NS1, Theorem 3.7] that there exists an injective map $\iota_N : \mathcal{B}_0(\lambda) \hookrightarrow \mathcal{B}_0(N\lambda)$ such that $\iota_N(u_\lambda) = u_{N\lambda}$, and

$$\begin{aligned} \text{wt}(\iota_N(b)) &= N \text{wt}(b), & \varepsilon_i(\iota_N(b)) &= N \varepsilon_i(b), & \varphi_i(\iota_N(b)) &= N \varphi_i(b), \\ \iota_N(e_i b) &= e_i^N \iota_N(b), & \iota_N(f_i b) &= f_i^N \iota_N(b), \end{aligned}$$

for $b \in \mathcal{B}_0(\lambda)$ and $i \in I_{\text{af}}$, where we understand that $\iota_N(\mathbf{0}) = \mathbf{0}$. Write λ as $\lambda = \sum_{i \in I} m_i \varpi_i$, with $m_i \in \mathbb{Z}_{\geq 0}$, $i \in I_0$. We deduce from [BN, Remark 4.17] (see also [Kas5, §13]) that there exists an embedding $\mathcal{B}_0(N\lambda) \hookrightarrow \tilde{\mathcal{B}} := \bigotimes_{i \in I} \mathcal{B}(\varpi_i)^{\otimes N m_i}$ of crystals that maps $u_{N\lambda}$ to $\tilde{u} := \bigotimes_{i \in I} u_{\varpi_i}^{\otimes N m_i}$. Also, we know from [Kas5, §10] that for every $i, j \in I$, there exists an isomorphism $\mathcal{B}(\varpi_i) \otimes \mathcal{B}(\varpi_j) \xrightarrow{\sim} \mathcal{B}(\varpi_j) \otimes \mathcal{B}(\varpi_i)$ of crystals that maps $u_{\varpi_i} \otimes u_{\varpi_j}$ to $u_{\varpi_j} \otimes u_{\varpi_i}$. Therefore, we obtain an isomorphism $\tilde{\mathcal{B}} \xrightarrow{\sim} \hat{\mathcal{B}} := \left(\bigotimes_{i \in I} \mathcal{B}(\varpi_i)^{\otimes m_i} \right)^{\otimes N}$ of crystals that maps $\tilde{u}$ to $\hat{u} := \left(\bigotimes_{i \in I} u_{\varpi_i}^{\otimes m_i} \right)^{\otimes N}$.

Combining the above, we see that $\mathcal{B}_0(N\lambda)$ is isomorphic, as a crystal, to the connected component of $\widehat{\mathcal{B}}$ containing $\widehat{u}$.

From [BN, Remark 4.17], we deduce that there exists an embedding $\mathcal{B}_0(\lambda)^{\otimes N} \hookrightarrow \widehat{\mathcal{B}}$ of crystals that maps $u_\lambda^{\otimes N}$ to $\widehat{u}$; note that the connected component of $\widehat{\mathcal{B}}$ containing $\widehat{u}$ (which is isomorphic as a crystal to $\mathcal{B}_0(N\lambda)$) is contained in the image of the embedding $\mathcal{B}_0(\lambda)^{\otimes N} \hookrightarrow \widehat{\mathcal{B}}$ of crystals. Hence we obtain an embedding $\iota' : \mathcal{B}_0(N\lambda) \hookrightarrow \mathcal{B}_0(\lambda)^{\otimes N}$ that maps $u_{N\lambda}$ to $u_\lambda^{\otimes N}$. Now it is clear that the composite $\iota' \circ \iota_N : \mathcal{B}_0(\lambda) \hookrightarrow \mathcal{B}_0(\lambda)^{\otimes N}$ satisfies conditions (5.2.1) and (5.2.2) required of σ_N . The uniqueness follows from the connectedness of $\mathcal{B}_0(\lambda)$. This proves the proposition. $\square$

Because $\sigma_{MN} = \sigma_M^{\otimes N} \circ \sigma_N$ for all $M, N \in \mathbb{Z}_{>0}$, we can prove the following proposition in exactly the same way as [Kas4, Proposition 8.3.2 (3)] and [NS1, Proposition 3.12].

Proposition 5.2.2. *Let $b \in \mathcal{B}_0(\lambda)$. There exists $N_b \in \mathbb{Z}_{>0}$ such that for every multiple $N \in \mathbb{Z}_{>0}$ of N_b , the element $\sigma_N(b) \in \mathcal{B}(\lambda)^{\otimes N}$ is of the form $\sigma_N(b) = u_{x_1} \otimes u_{x_2} \otimes \cdots \otimes u_{x_N}$ for some $x_1, x_2, \dots, x_N \in (W^J)_{\text{af}}$.*

Since $\langle c, \lambda \rangle = 0$, we see that $\{\langle \beta^\vee, \lambda \rangle \mid \beta \in \Delta_{\text{af}}\} = \{\langle \alpha^\vee, \lambda \rangle \mid \alpha \in \Delta\}$ is a finite set. Define $N_\lambda \in \mathbb{Z}_{>0}$ to be the least common multiple of the integers in the finite set $\{\langle \beta^\vee, \lambda \rangle \mid \beta \in \Delta_{\text{af}}^+\} \setminus \{0\}$.

Lemma 5.2.3. *Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ . If $\eta = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, then $Na_u \in \mathbb{Z}$ for all $0 \leq u \leq s$.*

Proof. If $u = 0$ or s , then the assertion is obvious. Assume that $1 \leq u \leq s - 1$. By the definition of a SiLS path, there exists a directed path from x_{u+1} to x_u in $\text{SiB}(\lambda; a_u)$; in particular, there exist $x, y \in (W^J)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$ such that $x \xrightarrow{\beta} y$ in $\text{SiB}(\lambda; a_u)$. By the definition of N_λ , it suffices to show that $a_u \langle x^{-1}\beta^\vee, \lambda \rangle \in \mathbb{Z} \setminus \{0\}$. But, since $x \xrightarrow{\beta} y$ is contained in $\text{SiB}(\lambda; a_u)$, we have $a_u \langle x^{-1}\beta^\vee, \lambda \rangle = a_u \langle \beta^\vee, x\lambda \rangle \in \mathbb{Z}$ by the definition. Suppose now that $\langle x^{-1}\beta^\vee, \lambda \rangle = 0$. Then, $x^{-1}\beta \in (\Delta_J)_{\text{af}}$, and hence $r_{x^{-1}\beta} \in (W_J)_{\text{af}}$ by Remark 2.3.1. Therefore, $y = r_\beta x = x r_{x^{-1}\beta} \notin (W^J)_{\text{af}}$, which is a contradiction. This proves the lemma. $\square$

Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ . We define $\sigma_N : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \hookrightarrow \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ as follows. Let $\eta = (x_1, \dots, x_s; a_0, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$. Note that $k_u := Na_u \in \mathbb{Z}$ for all $0 \leq u \leq s$ by Lemma 5.2.3. Now we set

$$\sigma_N(\eta) := \underbrace{\eta_{x_1} \otimes \cdots \otimes \eta_{x_1}}_{(k_1 - k_0) \text{ times}} \otimes \underbrace{\eta_{x_2} \otimes \cdots \otimes \eta_{x_2}}_{(k_2 - k_0) \text{ times}} \otimes \cdots \otimes \underbrace{\eta_{x_s} \otimes \cdots \otimes \eta_{x_s}}_{(k_s - k_{s-1}) \text{ times}} \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}.$$

Proposition 5.2.4. *Keep the notation and setting above. The map $\sigma_N : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \hookrightarrow \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ above is an injective map such that $\sigma_N(\eta_e) = \eta_e^{\otimes N}$, and*

$$\text{wt}(\sigma_N(\eta)) = N \text{wt}(\eta), \quad \varepsilon_i(\sigma_N(\eta)) = N \varepsilon_i(\eta), \quad \varphi_i(\sigma_N(\eta)) = N \varphi_i(\eta), \quad (5.2.3)$$

$$\sigma_N(e_i\eta) = e_i^N \sigma_N(\eta), \quad \sigma_N(f_i\eta) = f_i^N \sigma_N(\eta), \quad (5.2.4)$$

for all $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$, where we understand that $\sigma_N(\mathbf{0}) = \mathbf{0}$.

Proof. First, we define a map $\mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N} \rightarrow \mathbb{B}(\lambda)^{\otimes N}$ by $\eta_1 \otimes \cdots \otimes \eta_N \mapsto \overline{\eta_1} \otimes \cdots \otimes \overline{\eta_N}$; by abuse of notation, we also denote this map by $\overline{} : \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N} \rightarrow \mathbb{B}(\lambda)^{\otimes N}$. We see from Remark 4.3.1 (1) and the tensor product rule for crystals that this map $\overline{} : \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N} \rightarrow \mathbb{B}(\lambda)^{\otimes N}$ is a strict morphism of crystals.

For $\pi_1, \dots, \pi_N \in \mathbb{B}(\lambda)$, define the concatenation $\pi_1 * \cdots * \pi_N$ by

$$(\pi_1 * \cdots * \pi_N)(t) := \sum_{L=1}^{K-1} \pi_L(1) + \pi_K(Nt - K + 1) \\ \text{for } \frac{K-1}{N} \leq t \leq \frac{K}{N} \text{ and } 1 \leq K \leq N,$$

and set $\mathbb{B}(\lambda)^{*N} := \{\pi_1 * \cdots * \pi_N \mid \pi_K \in \mathbb{B}(\lambda), 1 \leq K \leq L\}$. For $\tilde{\pi} \in \mathbb{B}(\lambda)$, we set $\text{wt}(\tilde{\pi}) := \tilde{\pi}(1)$. Also, for $\tilde{\pi} \in \mathbb{B}(\lambda)$ and $i \in I_{\text{af}}$, we define $e_i \tilde{\pi}$ and $f_i \tilde{\pi}$ in exactly the same way as in Definition 2.2.6; we deduce from [Li2, Lemma 2.7] that $e_i \tilde{\pi}, f_i \tilde{\pi} \in \mathbb{B}(\lambda)^{*N} \cup \{\mathbf{0}\}$. Now, for $\tilde{\pi} \in \mathbb{B}(\lambda)$ and $i \in I_{\text{af}}$, define $\varepsilon_i(\tilde{\pi})$ and $\varphi_i(\tilde{\pi})$ as in (2.2.6). Then we deduce from [Li2, Lemma 2.7] that the set $\mathbb{B}(\lambda)^{*N}$, equipped with the maps $\text{wt}, e_i, f_i, i \in I$, and $\varepsilon_i, \varphi_i, i \in I$, is a crystal with weights in P_{af} , and that the map $\kappa : \mathbb{B}(\lambda)^{\otimes N} \rightarrow \mathbb{B}(\lambda)^{*N}$, $\pi_1 \otimes \cdots \otimes \pi_N \mapsto \pi_1 * \cdots * \pi_N$, is an isomorphism of crystals.

For $\pi \in \mathbb{B}(\lambda)$, we define $\iota_N(\pi)$ by $(\iota_N(\pi))(t) := N\pi(t)$ for $t \in [0, 1]$; it is easily seen that $N\pi \in \mathbb{B}(N\lambda)$. Thus we obtain an injective map $\iota_N : \mathbb{B}(\lambda) \hookrightarrow \mathbb{B}(N\lambda)$. We know from [Li2, Lemma 2.4] that

$$\begin{aligned} \text{wt}(\iota_N(\pi)) &= N \text{wt}(\pi), & \varepsilon_i(\iota_N(\pi)) &= N \varepsilon_i(\pi), & \varphi_i(\iota_N(\pi)) &= N \varphi_i(\pi), \\ \iota_N(e_i \pi) &= e_i^N \iota_N(\pi), & \iota_N(f_i \pi) &= f_i^N \iota_N(\pi), \end{aligned}$$

for $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$, where we understand that $\iota_N(\mathbf{0}) = \mathbf{0}$.

Observe that for every $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, the element $\iota_N(\overline{\eta}) \in \mathbb{B}(N\lambda)$ is identical to the element $\kappa(\overline{\sigma_N(\eta)}) \in \mathbb{B}(\lambda)^{*N}$ (as a piecewise-linear, continuous map from $[0, 1]$ to $\mathbb{R} \otimes_{\mathbb{Z}} P$). The equalities in (5.2.3) follow immediately from this fact and Remark 4.3.1 (2). Let us show the equalities in (5.2.4). We give a proof only for e_i , $i \in I_{\text{af}}$; the proof for f_i , $i \in I_{\text{af}}$, is similar. Let $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$ be such that $e_i \eta \neq \mathbf{0}$; note that $e_i \overline{\eta} \neq \mathbf{0}$, and $e_i^N \sigma_N(\eta) \neq \mathbf{0}$. Write $\sigma_N(\eta) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ as $\sigma_N(\eta) = \eta_{y_1} \otimes \cdots \otimes \eta_{y_N}$ with $y_1, \dots, y_N \in (W^J)_{\text{af}}$, and assume that

$$e_i^N \sigma_N(\eta) = e_i^N (\eta_{y_1} \otimes \cdots \otimes \eta_{y_N}) = e_i^{p_1} \eta_{y_1} \otimes \cdots \otimes e_i^{p_N} \eta_{y_N} \quad (5.2.5)$$

for some $p_1, \dots, p_N \in \mathbb{Z}_{\geq 0}$ such that $p_1 + \cdots + p_N = N$. Then we see that

$$e_i^N \kappa(\overline{\sigma_N(\eta)}) = \kappa(\overline{e_i^N \sigma_N(\eta)}) = e_i^{p_1} \overline{\eta_{y_1}} * \cdots * e_i^{p_N} \overline{\eta_{y_N}}. \quad (5.2.6)$$

Next, let us define $t_0, t_1 \in [0, 1]$ by (2.2.4), with $\pi = \bar{\eta}$. It follows from Lemma 5.2.3 and the definition of the root operator e_i that $t_0 = L/N$ and $t_1 = K/N$ for some $0 \leq L < K \leq N$; note that $\langle \alpha_i^\vee, y_M \lambda \rangle < 0$ for all $L+1 \leq M \leq K$ since the function $H_i^{\bar{\eta}}(t)$ is strictly decreasing on $[t_0, t_1]$. It is easily seen from the definition of the root operator e_i that $\sigma_N(e_i \eta) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ is of the form:

$$\sigma_N(e_i \eta) = \eta_{y_1} * \cdots * \eta_{y_L} * \underbrace{\eta_{r_i y_{L+1}} * \cdots * \eta_{r_i y_K}}_{\text{"reflected" by } r_i} * \eta_{y_{K+1}} * \cdots * \eta_{y_N}, \quad (5.2.7)$$

and hence $\kappa(\overline{\sigma_N(e_i \eta)}) \in \mathbb{B}(\lambda)^{*N}$ is of the form:

$$\kappa(\overline{\sigma_N(e_i \eta)}) = \overline{\eta_{y_1}} * \cdots * \overline{\eta_{y_L}} * \underbrace{\overline{\eta_{r_i y_{L+1}}} * \cdots * \overline{\eta_{r_i y_K}}}_{\text{"reflected" by } r_i} * \overline{\eta_{y_{K+1}}} * \cdots * \overline{\eta_{y_N}}. \quad (5.2.8)$$

Here, we have

$$\kappa(\overline{\sigma_N(e_i \eta)}) = \iota_N(\overline{e_i \eta}) = \iota_N(e_i \bar{\eta}) = e_i^N \iota_N(\bar{\eta}) = e_i^N \kappa(\overline{\sigma_N(\bar{\eta})}).$$

Combining this with (5.2.6) and (5.2.8), we obtain

$$\overline{\eta_{y_1}} * \cdots * \overline{\eta_{y_{L-1}}} * \underbrace{\overline{\eta_{r_i y_L}} * \cdots * \overline{\eta_{r_i y_K}}}_{\text{"reflected" by } r_i} * \overline{\eta_{y_{K+1}}} * \cdots * \overline{\eta_{y_N}} = e_i^{p_1} \overline{\eta_{y_1}} * \cdots * e_i^{p_N} \overline{\eta_{y_N}}.$$

Therefore, if we set $n_M := \langle \alpha_i^\vee, y_M \lambda \rangle < 0$ for $L+1 \leq M \leq K$, then

$$p_M = \begin{cases} -n_M & \text{if } L+1 \leq M \leq K, \\ 0 & \text{otherwise,} \end{cases}$$

for $1 \leq M \leq N$. From this and (5.2.5), we deduce that

$$\begin{aligned} e_i^N \sigma_N(\eta) &= \eta_{y_1} \otimes \cdots \otimes \eta_{y_L} \otimes e_i^{-n_{L+1}} \eta_{y_{L+1}} \otimes \cdots \otimes e_i^{-n_K} \eta_{y_K} \otimes \eta_{y_{K+1}} \otimes \cdots \otimes \eta_{y_N} \\ &= \eta_{y_1} \otimes \cdots \otimes \eta_{y_L} \otimes \eta_{r_i y_{L+1}} \otimes \cdots \otimes \eta_{r_i y_K} \otimes \eta_{y_{K+1}} \otimes \cdots \otimes \eta_{y_N} \quad \text{by (5.1.6)} \\ &= \sigma_N(e_i \eta) \quad \text{by (5.2.7).} \end{aligned}$$

This completes the proof of the proposition. $\square$

5.3 Proof of Proposition 3.2.2.

Lemma 5.3.1. *Let X be a monomial in the Kashiwara operators, i.e., $X = g_m g_{m-1} \cdots g_2 g_1$ with $g_k \in \{e_i, f_i \mid i \in I_{\text{af}}\}$ for $1 \leq k \leq m$.*

- (1) *If $X u_\lambda \neq \mathbf{0}$, then $X \eta_e \neq \mathbf{0}$. We set $b_k := g_k g_{k-1} \cdots g_1 u_\lambda$ for $0 \leq k \leq m$, and let $N \in \mathbb{Z}_{>0}$ be a common multiple of N_{b_k} , $0 \leq k \leq m$, and N_λ (see Proposition 5.2.2 and the comment preceding Lemma 5.2.3). If $\sigma_N(X u_\lambda) = u_{x_1} \otimes \cdots \otimes u_{x_N}$ with $x_1, \dots, x_N \in (W^J)_{\text{af}}$, then $\sigma_N(X \eta_e) = \eta_{x_1} \otimes \cdots \otimes \eta_{x_N}$.*

- (2) If $X\eta_e \neq \mathbf{0}$, then $Xu_\lambda \neq \mathbf{0}$. Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ , and write $\sigma_N(X\eta_e)$ as $\sigma_N(X\eta_e) = \eta_{x_1} \otimes \cdots \otimes \eta_{x_N}$ for some $x_1, \dots, x_N \in (W^J)_{\text{af}}$. Then, $\sigma_N(Xu_\lambda) = u_{x_1} \otimes \cdots \otimes u_{x_N}$.

Proof. We give a proof only for part (1); the proof for part (2) is similar. We proceed by induction on m . If $m = 0$, then the assertion is obvious. Assume that $m > 0$, and take $i \in I_{\text{af}}$ for which $g_m = e_i$ or f_i . Set $Y := g_{m-1} \cdots g_2 g_1$, and write $\sigma_N(Yu_\lambda) = \sigma_N(b_{m-1})$ as $\sigma_N(Yu_\lambda) = u_{y_1} \otimes \cdots \otimes u_{y_N}$ for some $y_1, \dots, y_N \in (W^J)_{\text{af}}$. Then, by our induction hypothesis, $Y\eta_e \neq \mathbf{0}$, and $\sigma_N(Y\eta_e) = \eta_{y_1} \otimes \cdots \otimes \eta_{y_N}$. Also, we see from the tensor product rule for crystals, using (5.1.5), that $\varepsilon_i(\sigma_N(Yu_\lambda)) = \varepsilon_i(\sigma_N(Y\eta_e))$ and $\varphi_i(\sigma_N(Yu_\lambda)) = \varphi_i(\sigma_N(Y\eta_e))$, and hence that $\varepsilon_i(Yu_\lambda) = \varepsilon_i(Y\eta_e)$ and $\varphi_i(Yu_\lambda) = \varphi_i(Y\eta_e)$ by (5.2.1), (5.2.3). Therefore, $Xu_\lambda = g_m Yu_\lambda \neq \mathbf{0}$ implies that $X\eta_e = g_m Y\eta_e \neq \mathbf{0}$. Furthermore, if

$$\sigma_N(Xu_\lambda) = g_m^N(u_{y_1} \otimes \cdots \otimes u_{y_N}) = g_m^{p_1} u_{y_1} \otimes \cdots \otimes g_m^{p_N} u_{y_N}$$

for some $p_1, \dots, p_N \in \mathbb{Z}_{\geq 0}$ with $p_1 + \cdots + p_N = N$, then it follows from the tensor product rule for crystals, together with (5.1.5), that

$$\sigma_N(X\eta_e) = g_m^N(\eta_{y_1} \otimes \cdots \otimes \eta_{y_N}) = g_m^{p_1} \eta_{y_1} \otimes \cdots \otimes g_m^{p_N} \eta_{y_N}.$$

Since $g_m^{p_1} u_{y_1} \otimes \cdots \otimes g_m^{p_N} u_{y_N} = \sigma_N(Xu_\lambda) = u_{x_1} \otimes \cdots \otimes u_{x_N}$ by the assumption, we deduce, by using (5.1.6), that $g_m^{p_L} \eta_{y_L} = \eta_{x_L}$ for all $1 \leq L \leq N$. This proves part (1). $\square$

Proof of Proposition 3.2.2. It suffices to show the following for monomials X, Y in the Kashiwara operators (cf. [Kas3, Proof of Theorem 4.1] and [NS1, Proof of Theorem 5.1]):

- (i) $Xu_\lambda \neq \mathbf{0}$ in $\mathcal{B}_0(\lambda)$ if and only if $X\eta_e \neq \mathbf{0}$ in $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$;
- (ii) $Xu_\lambda = Yu_\lambda$ in $\mathcal{B}_0(\lambda)$ if and only if $X\eta_e = Y\eta_e$ in $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$.

Part (i) has already been shown in Lemma 5.3.1. Let us show part (ii). Assume that $Xu_\lambda = Yu_\lambda \neq \mathbf{0}$. By Lemma 5.3.1 (1), we have $X\eta_e \neq \mathbf{0}$ and $Y\eta_e \neq \mathbf{0}$. Take a multiple $N \in \mathbb{Z}_{>0}$ of N_λ such that the assumption of Lemma 5.3.1 (1) is satisfied for both of the elements Xu_λ and Yu_λ , and write $\sigma_N(Xu_\lambda)$ and $\sigma_N(Yu_\lambda)$ as:

$$\sigma_N(Xu_\lambda) = u_{x_1} \otimes u_{x_2} \otimes \cdots \otimes u_{x_N}, \quad \sigma_N(Yu_\lambda) = u_{y_1} \otimes u_{y_2} \otimes \cdots \otimes u_{y_N},$$

for some $x_1, \dots, x_N \in (W^J)_{\text{af}}$ and $y_1, \dots, y_N \in (W^J)_{\text{af}}$. Then, by Lemma 5.3.1 (1),

$$\sigma_N(X\eta_e) = \eta_{x_1} \otimes \eta_{x_2} \otimes \cdots \otimes \eta_{x_N}, \quad \sigma_N(Y\eta_e) = \eta_{y_1} \otimes \eta_{y_2} \otimes \cdots \otimes \eta_{y_N}.$$

Since $Xu_\lambda = Yu_\lambda$, we have $x_L = y_L$ for all $1 \leq L \leq N$. Therefore, we deduce that $\sigma_N(X\eta_e) = \sigma_N(Y\eta_e)$, and hence $X\eta_e = Y\eta_e$ by the injectivity of σ_N . Thus, we have proved the “only if” part of part (ii). The “if” part can be shown similarly; use Lemma 5.3.1 (2) instead of Lemma 5.3.1 (1). This completes the proof of Proposition 3.2.2. $\square$

Remark 5.3.2. Since $\mathcal{B}_0(\lambda) \cong \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ as crystals, we can define the action of the Weyl group W_{af} on $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ by the same formula as (5.1.1) for the one on $\mathcal{B}_0(\lambda)$. Moreover, from equation (5.1.6), we see that the element $\eta_e = (e; 0, 1) \in \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ is an extremal element.

6 Directed paths in $\text{SiB}(\lambda; a)$.

6.1 Some technical lemmas.

Let J, K be subsets of I such that $K \subset J$; we see by the definitions that $(W^J)_{\text{af}} \subset (W^K)_{\text{af}}$.

Lemma 6.1.1. *Let J, K be subsets of I such that $K \subset J$, and let $x \in (W^K)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$ be such that $x \xrightarrow{\beta} r_{\beta}x$ in SiB^K . Then there exists a directed path from $\Pi^J(x)$ to $\Pi^J(r_{\beta}x)$ in SiB^J .*

Proof. Let λ and Λ be (arbitrary) elements in P^+ such that $J_{\lambda} = \{i \in I \mid \langle \alpha_i^{\vee}, \lambda \rangle = 0\} = J$, and $J_{\Lambda} = \{i \in I \mid \langle \alpha_i^{\vee}, \Lambda \rangle = 0\} = K$.

Claim 1. *If β is a simple root, then the assertion of the lemma holds.*

Proof of Claim 1. Assume that $\beta = \alpha_i$ for some $i \in I_{\text{af}}$. Because $x \xrightarrow{\alpha_i} r_i x$ in SiB^K , we see from (4.1.2) that $\langle \alpha_i^{\vee}, x\Lambda \rangle > 0$, and hence $\langle \alpha_i^{\vee}, x\lambda \rangle \geq 0$ since $K \subset J$. Assume that $\langle \alpha_i^{\vee}, x\lambda \rangle > 0$. Since $x\lambda = \Pi^J(x)\lambda$, we see from Lemma 2.3.6 and (4.1.2) that $r_i \Pi^J(x) \in (W^J)_{\text{af}}$, and $\Pi^J(x) \xrightarrow{\alpha_i} r_i \Pi^J(x)$ in SiB^J ; note that $r_i \Pi^J(x) = \Pi^J(r_i \Pi^J(x)) = \Pi^J(r_i x)$. Thus we obtain $\Pi^J(x) \xrightarrow{\alpha_i} \Pi^J(r_i x)$ in SiB^J . If $\langle \alpha_i^{\vee}, x\lambda \rangle = 0$, then we see that $x^{-1}\alpha_i \in (\Delta_J)_{\text{af}}$, which implies that $\Pi^J(r_i x) = \Pi^J(xr_{x^{-1}\alpha_i}) = \Pi^J(x)$. This proves Claim 1. ■

Let us consider the case of general $\beta \in \Delta_{\text{af}}^+$. By [AK, Lemma 1.4], there exist $i_1, i_2, \dots, i_k \in I_{\text{af}}$ such that

$$x\Lambda = \mu_k \xrightarrow{\alpha_{i_k}} \mu_{k-1} \xrightarrow{\alpha_{i_{k-1}}} \dots \xrightarrow{\alpha_{i_2}} \mu_1 \xrightarrow{\alpha_{i_1}} \mu_0,$$

with $\mu_0 \equiv \Lambda \pmod{\mathbb{Q}\delta}$, where $\mu_m = r_{i_{m+1}} \dots r_{i_k} x\Lambda$ for $0 \leq m \leq k$. We show the assertion by induction on k . Assume that $k = 0$; observe that in this case, $x = z_{\xi} t_{\xi}$ for some $\xi \in Q^{\vee, K\text{-ad}}$ since $x\Lambda \equiv \Lambda \pmod{\mathbb{Q}\delta}$. It follows from Lemma 4.1.7 that $\beta = \alpha_i$ for some $i \in I \setminus K$, and hence the assertion follows immediately from Claim 1. Assume that $k > 0$. For simplicity of notation, we set $y := r_{\beta}x \in (W^K)_{\text{af}}$ and $i := i_k$. Then, by (4.1.2), we have $x \xrightarrow{\alpha_i} r_i x$ in SiB^K . Write $y^{-1}\alpha_i = \alpha + n\delta$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}$; by Lemma 4.1.4, $\alpha \notin \Delta_K^+$. If α is a negative root, then $\beta = \alpha_i$ by Lemma 4.1.4, and hence the assertion follows immediately from Claim 1. Assume that $\alpha \in \Delta^+ \setminus \Delta_K^+$. By Lemma 4.1.4 applied to the subset K of I , we obtain the following diagram in SiB^K :

$$\begin{array}{ccc} x & \xrightarrow{\beta} & y \\ \alpha_i \downarrow & & \vdots \downarrow \alpha_i \\ r_i x & \xrightarrow{\dots \dots r_i \beta} & r_i y \end{array} \tag{6.1.1}$$

We see from Claim 1 that there exists a directed path from $\Pi^J(x)$ to $\Pi^J(r_i x)$ in SiB^J . Also, by our induction hypothesis applied to $r_i x$, there exists a directed path from $\Pi^J(r_i x)$ to $\Pi^J(r_i y)$ in SiB^J . Concatenating these, we obtain a directed path from $\Pi^J(x)$ to $\Pi^J(r_i y)$ in SiB^J , which proves the lemma in the case that $\Pi^J(r_i y) = \Pi^J(y)$. Now, let us assume that $\Pi^J(r_i y) \neq \Pi^J(y)$. We see from (6.1.1) and (4.1.2) that $\langle \alpha_i^\vee, y\lambda \rangle > 0$, and hence $\langle \alpha_i^\vee, y\lambda \rangle \geq 0$. Since $\Pi^J(yr_{y^{-1}\alpha_i}) = \Pi^J(r_i y) \neq \Pi^J(y)$ by our assumption, it follows from Proposition 2.3.2, together with Remark 2.3.1, that $\langle \alpha_i^\vee, y\lambda \rangle \neq 0$, and hence $\langle \alpha_i^\vee, y\lambda \rangle > 0$. Since $\Pi^J(y)\lambda = y\lambda$, we see from Lemma 2.3.6 that $r_i \Pi^J(y) \in (W^J)_{\text{af}}$; note that $r_i \Pi^J(y) = \Pi^J(r_i \Pi^J(y)) = \Pi^J(r_i y)$, and $\langle \alpha_i^\vee, \Pi^J(r_i y)\lambda \rangle < 0$. Similarly, we can show that $\langle \alpha_i^\vee, \Pi^J(x)\lambda \rangle \geq 0$ since $x \xrightarrow{\alpha_i} r_i x$ in SiB^K . Therefore, by applying Lemma 4.1.6 (3) to the directed path from $\Pi^J(x)$ to $\Pi^J(r_i y)$ in SiB^J obtained above, we see that there exists a directed path from $\Pi^J(x)$ to $r_i \Pi^J(r_i y) = \Pi^J(y)$ in SiB^J . This proves the lemma. $\square$

Here, we make the following remark.

Remark 6.1.2. Let J be a subset of I . Let $x, y \in (W^J)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$ be such that $x \xrightarrow{\beta} y$ in SiB^J . Write x and y as $x = wz_\xi t_\xi$ and $y = vz_\zeta t_\zeta$, with $w, v \in W$ and $\xi, \zeta \in Q^{\vee, J\text{-ad}}$ (see (2.3.5)). We see from Corollary 4.2.2 that $\beta = w\gamma + n\delta$ for some $\gamma \in \Delta^+ \setminus \Delta_J^+$ and $n \in \{0, 1\}$. Also, it follows easily that $\zeta - \xi = nz_\xi^{-1}\gamma^\vee$. In particular, we have $[\zeta - \xi] = n[\gamma^\vee] \in Q_{J^c}^{\vee+} = \sum_{j \in J^c} \mathbb{Z}_{\geq 0} \alpha_j^\vee$, where $J^c := I \setminus J$, and $[\cdot] = [\cdot]_{J^c} : Q^\vee \rightarrow Q_{J^c}^\vee = \bigoplus_{j \in J^c} \mathbb{Z} \alpha_j^\vee$ denote the projection from $Q^\vee = Q_{J^c}^\vee \oplus Q_J^\vee$ onto $Q_{J^c}^\vee$ with kernel $Q_J^\vee$.

Lemma 6.1.3. *Let J be a subset of I . For each $i \in J^c = I \setminus J$ and $\xi \in Q^{\vee, J\text{-ad}}$, there exists a positive real root $\beta \in \Delta_{\text{af}}^+$ of the form $\beta = -\gamma + \delta$, with $\gamma \in \Delta^+ \setminus \Delta_J^+$, satisfying the conditions that $[\gamma^\vee] = \alpha_i^\vee$, $r_\beta z_\xi t_\xi \in (W^J)_{\text{af}}$, and $r_\beta z_\xi t_\xi \xrightarrow{\beta} z_\xi t_\xi$ in SiB^J .*

Proof. Let $i \in I \setminus J$, and let $\xi \in Q^{\vee, J\text{-ad}}$. Since $\ell^{\frac{\infty}{2}}(r_i t_{\xi - \alpha_i^\vee}) = \ell(r_i) + 2\langle \xi - \alpha_i^\vee, \rho \rangle = 2\langle \xi, \rho \rangle - 1 = \ell^{\frac{\infty}{2}}(t_\xi) - 1$, we see that $r_i t_{\xi - \alpha_i^\vee} = r_{\beta'} t_\xi \xrightarrow{\beta'} t_\xi$ in $\text{SiB}^\emptyset$, with $\beta' = -\alpha_i + \delta$. Therefore, by Lemma 6.1.1, there exists a directed path from $\Pi^J(r_i t_{\xi - \alpha_i^\vee})$ to $\Pi^J(t_\xi)$ in SiB^J . Since $i \in I \setminus J$, we see from Lemma 2.3.5 (2) that $\Pi^J(r_i t_{\xi - \alpha_i^\vee}) = r_i z_{\xi - \alpha_i^\vee} t_{\zeta'}$, with $\zeta' = \xi - \alpha_i^\vee + \phi_J(\xi - \alpha_i^\vee)$. Also, we see from Lemma 2.3.5 (1), (2) that $\Pi^J(t_\xi) = z_\xi t_\xi$. Remark that $[\xi - \zeta'] = \alpha_i^\vee$. Let $x \xrightarrow{\beta} y := z_\xi t_\xi$, $x \in (W^J)_{\text{af}}$, $\beta \in \Delta_{\text{af}}^+$, be the final edge of the directed path in SiB^J from $\Pi^J(r_i t_{\xi - \alpha_i^\vee})$ to $\Pi^J(t_\xi) = z_\xi t_\xi$ above:

$$\Pi^J(r_i t_{\xi - \alpha_i^\vee}) \rightarrow \cdots \rightarrow x \xrightarrow{\beta} y = z_\xi t_\xi \quad \text{in } \text{SiB}^J. \quad (6.1.2)$$

Here, let us write x as $x = vz_\zeta t_\zeta$, with $v \in W^J$ and $\zeta \in Q^{\vee, J\text{-ad}}$. By applying Remark 6.1.2 to each edge of the directed path from $\Pi^J(r_i t_{\xi - \alpha_i^\vee}) = r_i z_{\xi - \alpha_i^\vee} t_{\zeta'}$ to $x = vz_\zeta t_\zeta$ (resp., the final edge $x \xrightarrow{\beta} y = z_\xi t_\xi$), we obtain $[\zeta - \zeta'] \in Q_{J^c}^{\vee+}$ (resp., $[\xi - \zeta] \in Q_{J^c}^{\vee+}$). Therefore, we deduce

that

$$\alpha_i^\vee = [\xi - \zeta'] = \underbrace{[\xi - \zeta]}_{\in Q_{J^c}^{\vee+}} + \underbrace{[\zeta - \zeta']}_{\in Q_J^{\vee+}},$$

which implies that $[\xi - \zeta] = 0$ or $\alpha_i^\vee$. Now, let $\lambda \in P^+$ be an (arbitrary) element such that $J_\lambda = J$. It follows immediately from Proposition 4.2.1 that $x\lambda \xrightarrow{\beta} y\lambda = z_\xi t_\xi \lambda$ in $\text{LZ}(\lambda)$. Since $z_\xi t_\xi \lambda \equiv \lambda \pmod{\mathbb{Q}\delta}$, we see that β is of the form $\beta = -\gamma + \delta$ for some $\gamma \in \Delta^+ \setminus \Delta_J^+$. Hence a direct computation shows that $\zeta = \xi - z_\xi^{-1} \gamma^\vee$; note that $[z_\xi^{-1} \gamma^\vee] = [\gamma^\vee] \neq 0$. Since $[\xi - \zeta] = 0$ or α_i as seen above, we conclude that $[\gamma^\vee] = \alpha_i$. This proves the lemma. $\square$

6.2 Directed paths from a translation to another translation.

In this subsection, we fix $\lambda \in P^+$, and set $J := J_\lambda = \{i \in I \mid \langle \alpha_i^\vee, \lambda \rangle = 0\}$. Recall that $[\cdot] = [\cdot]_{J^c} : Q^\vee \twoheadrightarrow Q_{J^c}^\vee \oplus Q_J^\vee$ denote the projection from $Q^\vee = Q_{J^c}^\vee \oplus Q_J^\vee$ onto $Q_{J^c}^\vee$ with kernel $Q_J^\vee$, where $J^c = I \setminus J$. Also, for a rational number $0 < a \leq 1$, we set

$$J^c(\lambda; a) := \{i \in J^c \mid a\langle \alpha_i^\vee, \lambda \rangle \in \mathbb{Z}\}, \quad I(\lambda; a) := \{i \in I \mid a\langle \alpha_i^\vee, \lambda \rangle \in \mathbb{Z}\} = J^c(\lambda; a) \cup J.$$

Lemma 6.2.1. *Let $\zeta, \xi \in Q^{\vee, J\text{-ad}}$, and let $0 < a \leq 1$ be a rational number. If $[\xi - \zeta] \in Q_{J^c(\lambda; a)}^{\vee+}$, then there exists a directed path from $z_\zeta t_\zeta$ to $z_\xi t_\xi$ in $\text{SiB}(\lambda; a)$.*

Proof. For $\alpha^\vee = \sum_{i \in I} c_i \alpha_i^\vee$ with $c_i \in \mathbb{Z}_{\geq 0}$, $i \in I$, we set $\text{ht}(\alpha^\vee) := \sum_{i \in I} c_i \in \mathbb{Z}_{\geq 0}$. We prove the lemma by induction on $\text{ht}([\xi - \zeta])$. Assume first that $\text{ht}([\xi - \zeta]) = 0$. Then, we have $[\xi - \zeta] = 0$, which implies that $\xi - \zeta \in Q_J^\vee$, and hence $\xi = \zeta$ by Lemma 2.3.5(1). Thus there exists a directed path (of length zero) from $z_\zeta t_\zeta$ to $z_\xi t_\xi = z_\zeta t_\zeta$ in $\text{SiB}(\lambda; a)$.

Assume next that $\text{ht}([\xi - \zeta]) > 0$. Take $i \in J^c(\lambda; a)$ such that $[\xi - \zeta] - \alpha_i^\vee \in Q_{J^c(\lambda; a)}^{\vee+}$, and set $\xi' := \xi - \alpha_i^\vee + \phi_J(\xi - \alpha_i^\vee) \in Q^{\vee, J\text{-ad}}$ (see Lemma 2.3.5(1)); note that $[\xi'] = [\xi] - \alpha_i^\vee$, and hence $\text{ht}([\xi' - \zeta]) = \text{ht}([\xi - \zeta]) - 1$. Therefore, by our induction hypothesis, there exists a directed path from $z_\zeta t_\zeta$ to $z_{\xi'} t_{\xi'}$ in $\text{SiB}(\lambda; a)$. Hence it suffices to show that there exists a directed path from $z_{\xi'} t_{\xi'}$ to $z_\xi t_\xi$ in $\text{SiB}(\lambda; a)$. We take $\beta = -\gamma + \delta$ of Lemma 6.1.3 (for the $\xi \in Q^{\vee, J\text{-ad}}$ and the $i \in J^c(\lambda; a)$ above). Since $[\gamma^\vee] = \alpha_i^\vee$ and $i \in J^c(\lambda; a)$, we see that the edge $r_\beta z_\xi t_\xi \xrightarrow{\beta} z_\xi t_\xi$ in SiB^J is in fact an edge in $\text{SiB}(\lambda; a)$. Write $r_\beta z_\xi t_\xi$ as $r_\beta z_\xi t_\xi = w z_{\xi''} t_{\xi''}$, with $w \in W^J$ and $\xi'' \in Q^{\vee, J\text{-ad}}$. Since $r_\beta z_\xi t_\xi = r_\gamma t_{-\gamma^\vee} z_\xi t_\xi$ by (2.1.2), we have $\xi'' = \xi - z_\xi^{-1} \gamma^\vee$. Because $[\xi' - \xi''] = ([\xi] - \alpha_i^\vee) - ([\xi] - [z_\xi^{-1} \gamma^\vee]) = ([\xi] - \alpha_i^\vee) - ([\xi] - \alpha_i^\vee) = 0$, we deduce from Lemma 2.3.5(1) that $\xi'' = \xi'$. Also, we see that $w \in W^J$ is equal to $r_\gamma z_\xi z_{\xi'}^{-1}$. Since $[\gamma^\vee] = \alpha_i^\vee$ and $i \in J^c(\lambda; a)$, it follows immediately that $\gamma \in \Delta_{I(\lambda; a)}$, and hence $r_\gamma \in W_{I(\lambda; a)} = \langle r_j \mid j \in I(\lambda; a) \rangle$. This implies that $w = r_\gamma z_\xi z_{\xi'}^{-1} \in W_{I(\lambda; a)}$ since $z_\xi z_{\xi'}^{-1} \in W_J \subset W_{I(\lambda; a)}$. Therefore, if $w = r_{i_1} r_{i_2} \cdots r_{i_k}$ is a reduced expression of w , then $i_l \in I(\lambda; a)$ for all $1 \leq l \leq k$. From this, it follows (see also the argument in Step 1 in the proof of Proposition 4.2.1) that

$$z_{\xi'} t_{\xi'} \xrightarrow{\alpha_{i_k}} r_{i_k} z_{\xi'} t_{\xi'} \xrightarrow{\alpha_{i_{k-1}}} r_{i_{k-1}} r_{i_k} z_{\xi'} t_{\xi'} \xrightarrow{\alpha_{i_{k-2}}} \cdots \xrightarrow{\alpha_{i_1}} r_{i_1} r_{i_2} \cdots r_{i_k} z_{\xi'} t_{\xi'} = w z_{\xi'} t_{\xi'}$$

is a directed path in $\text{SiB}(\lambda; a)$. Concatenating this directed path with $wz_{\xi'}t_{\xi'} = r_{\beta}z_{\xi}t_{\xi} \xrightarrow{\beta} z_{\xi}t_{\xi}$ above, we obtain a directed path from $z_{\xi'}t_{\xi'}$ to $z_{\xi}t_{\xi}$ in $\text{SiB}(\lambda; a)$, as desired. $\square$

Proposition 6.2.2. *Let $\zeta, \xi \in Q^{\vee, J\text{-ad}}$, and let $0 < a \leq 1$ be a rational number. There exists a directed path of nonzero length from $z_{\zeta}t_{\zeta}$ to $z_{\xi}t_{\xi}$ in $\text{SiB}(\lambda; a)$ if and only if the set $J^c(\lambda; a) = \{i \in J^c \mid a\langle\alpha_i^{\vee}, \lambda\rangle \in \mathbb{Z}\}$ is not empty, and $[\xi - \zeta] \in Q_{J^c(\lambda; a)}^{\vee+} \setminus \{0\}$.*

Proof. We prove the “only if” part. Let

$$z_{\zeta}t_{\zeta} = y_0 \xrightarrow{\beta_1} y_1 \xrightarrow{\beta_2} \cdots \xrightarrow{\beta_k} y_k = z_{\xi}t_{\xi}$$

be a directed path of nonzero length from $z_{\zeta}t_{\zeta}$ to $z_{\xi}t_{\xi}$ in $\text{SiB}(\lambda; a)$. By Lemma 4.1.7, we see that $\beta_1 = \alpha_i$ for some $i \in J^c$. Since $a\langle\alpha_i^{\vee}, \lambda\rangle = a\langle\beta_1^{\vee}, z_{\zeta}t_{\zeta}\lambda\rangle \in \mathbb{Z}$, it follows immediately that $i \in J^c(\lambda; a)$, and in particular, $J^c(\lambda; a) \neq \emptyset$. We write $y_l \in (W^J)_{\text{af}}$, $0 \leq l \leq k$, and $\beta_l \in \Delta_{\text{af}}^+$ (see Corollary 4.2.2), $1 \leq l \leq k$, as:

$$\begin{cases} y_l = w_l z_{\xi_l} t_{\xi_l} & \text{with } w_l \in W^J \text{ and } \xi_l \in Q^{\vee, J\text{-ad}}, \\ \beta_l = w_{l-1} \gamma_l + n_l \delta & \text{with } \gamma_l \in \Delta^+ \setminus \Delta_J^+ \text{ and } n_l \in \{0, 1\}. \end{cases}$$

Then we deduce from Remark 6.1.2 that

$$\xi - \zeta = \sum_{l=1}^k (\xi_l - \xi_{l-1}) = \sum_{l=1}^k n_l z_{\xi_{l-1}}^{-1} \gamma_l^{\vee}, \quad \text{and hence} \quad [\xi - \zeta] = \sum_{l=1}^k n_l [\gamma_l^{\vee}] \in Q_{J^c}^{\vee+}. \quad (6.2.1)$$

Also, using (2.3.5), we see by direct computation that $w_l = \lfloor w_{l-1} r_{\gamma_l} \rfloor$ for all $1 \leq l \leq k$. Let us show by induction on l that $w_l \in W_{I(\lambda; a)}$ and $\gamma_l \in \Delta_{I(\lambda; a)}$ for all $1 \leq l \leq k$. Since $w_0 = e$, and since $\beta_1 = \alpha_i$ with $i \in J^c(\lambda; a) \subset I(\lambda; a)$ as seen above, it follows immediately that $\gamma_1 = \alpha_i \in \Delta_{I(\lambda; a)}$ and $w_1 = \lfloor w_0 r_{\gamma_1} \rfloor = \lfloor r_i \rfloor \in W_{I(\lambda; a)}$. Assume that $l > 0$. By Proposition 4.2.1, we have $y_{l-1} \lambda \xrightarrow{\beta_l} y_l \lambda$ in $\text{LZ}(\lambda; a)$. We deduce from [NS4, Lemma 3.11], together with [NS4, Lemma 2.13], that $w_{l-1} \gamma_l \in \Delta_{I(\lambda; a)}$. Since $w_{l-1} \in W_{I(\lambda; a)}$ by our induction hypothesis, we obtain $\gamma_l \in \Delta_{I(\lambda; a)}$. Also, since $w_l = \lfloor w_{l-1} r_{\gamma_l} \rfloor$ with $w_{l-1} \in W_{I(\lambda; a)}$ and $\gamma_l \in \Delta_{I(\lambda; a)}$, it follows immediately that $w_l \in W_{I(\lambda; a)}$. Thus we have shown that $w_l \in W_{I(\lambda; a)}$ and $\gamma_l \in \Delta_{I(\lambda; a)}$ for all $1 \leq l \leq k$. Combining this fact and (6.2.1), we conclude that $[\xi - \zeta] \in Q_{J^c(\lambda; a)}^{\vee+}$. Suppose, for a contradiction, that $[\xi - \zeta] = 0$. Then, we have $\xi - \zeta \in Q_J^{\vee}$. This implies that $\xi = \zeta$ by Lemma 2.3.5(1), which contradicts the assumption. Thus we have proved the “only if” part. The “if” part follows immediately from Lemma 6.2.1. This completes the proof of the proposition. $\square$

7 Connected components of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$.

Throughout this section, we fix $\lambda \in P^+$, and set $J := J_{\lambda} = \{i \in I \mid \langle\alpha_i^{\vee}, \lambda\rangle = 0\}$. Write $\lambda \in P^+$ as $\lambda = \sum_{i \in I} m_i \varpi_i$ with $m_i \in \mathbb{Z}_{\geq 0}$, $i \in I$, and set

$$\text{Turn}(\lambda) := \{k/m_i \mid i \in J^c = I \setminus J, 0 \leq k \leq m_i\};$$

note that $m_i = \langle \alpha_i^\vee, \lambda \rangle$. Also, for simplicity of notation, we set $T_\xi := \Pi^J(t_\xi) = z_\xi t_\xi \in (W^J)_{\text{af}}$ for $\xi \in Q^{\vee, J\text{-ad}}$.

7.1 An extremal element in each connected component.

The next proposition follows immediately from Proposition 6.2.2.

Proposition 7.1.1. *Let $\xi_1, \dots, \xi_{s-1}, \xi_s \in Q^{\vee, J\text{-ad}}$. An element*

$$\eta = (T_{\xi_1}, \dots, T_{\xi_{s-1}}, T_{\xi_s}; a_0, a_1, \dots, a_{s-1}, a_s) \quad (7.1.1)$$

is contained in $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ if and only if $a_u \in \text{Turn}(\lambda)$ for all $0 \leq u \leq s$ and $[\xi_u - \xi_{u+1}] \in Q_{J^c(\lambda; a_u)}^{\vee+} \setminus \{0\}$ for all $1 \leq u \leq s-1$.

Proposition 7.1.2. *Each connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ contains a unique element of the form (7.1.1) with $\xi_s = 0$.*

In order to prove this proposition, we need some lemmas. Let $N \in \mathbb{Z}_{>0}$. For simplicity of notation, we set $[y_1, y_2, \dots, y_N] := \eta_{y_1} \otimes \eta_{y_2} \otimes \dots \otimes \eta_{y_N} \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ for $y_1, y_2, \dots, y_N \in (W^J)_{\text{af}}$.

Lemma 7.1.3. *Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ (see the comment preceding Lemma 5.2.3). Let $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, and write $\sigma_N(\eta) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ as $\sigma_N(\eta) = [y_1, y_2, \dots, y_N]$, with $y_1, y_2, \dots, y_N \in (W^J)_{\text{af}}$. Let X be a monomial in the root operators e_i and f_i , $i \in I_{\text{af}}$, and assume that $X\eta \neq \mathbf{0}$. Then, $\sigma_N(X\eta) = [v_1 y_1, v_2 y_2, \dots, v_N y_N]$ for some $v_1, v_2, \dots, v_N \in W_{\text{af}}$ such that $v_M y_M \in (W^J)_{\text{af}}$, $1 \leq M \leq N$.*

Proof. It suffices to show the assertion in the case when $X = e_i$ or f_i , $i \in I_{\text{af}}$; the assertion for a general X follows immediately by induction. From the definition of root operators, we obtain

$$\sigma_N(X\eta) = [y_1, \dots, y_L, r_i y_{L+1}, \dots, r_i y_K, y_{K+1}, \dots, y_N]$$

for some $1 \leq L < K \leq N$, with $r_i y_M \in (W^J)_{\text{af}}$ for all $L+1 \leq M \leq K$. This proves the lemma. $\square$

Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ . Let $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ be of the form (7.1.1). Then, $\sigma_N(\eta)$ is of the form $\sigma_N(\eta) = [T_{\zeta_1}, T_{\zeta_2}, \dots, T_{\zeta_N}]$ for some $\zeta_1, \zeta_2, \dots, \zeta_N \in Q^{\vee, J\text{-ad}}$. Let X be a monomial in the root operators e_i and f_i , $i \in I_{\text{af}}$, such that $X\eta \neq \mathbf{0}$; by Lemma 7.1.3, $\sigma_N(X\eta)$ is of the form $\sigma_N(X\eta) = [v_1 T_{\zeta_1}, v_2 T_{\zeta_2}, \dots, v_N T_{\zeta_N}]$ for some $v_1, v_2, \dots, v_N \in W_{\text{af}}$ such that $v_M T_{\zeta_M} = v_M z_{\zeta_M} t_{\zeta_M} \in (W^J)_{\text{af}}$, $1 \leq M \leq N$; note that $v_M \in (W^J)_{\text{af}}$ for all $1 \leq M \leq N$ by Lemma 2.3.7.

Lemma 7.1.4. *Keep the notation and setting above. Let $\eta' \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ be another element of the form (7.1.1), and write $\sigma_N(\eta')$ as $\sigma_N(\eta') = [T_{\zeta'_1}, T_{\zeta'_2}, \dots, T_{\zeta'_N}]$ for some $\zeta'_1, \zeta'_2, \dots, \zeta'_N \in Q^{\vee, J\text{-ad}}$. Then, $X\eta' \neq \mathbf{0}$, and $\sigma_N(X\eta') = [v_1 T_{\zeta'_1}, v_2 T_{\zeta'_2}, \dots, v_N T_{\zeta'_N}]$; note that $v_M T_{\zeta'_M} \in (W^J)_{\text{af}}$ for all $1 \leq M \leq N$ by Lemma 2.3.7 since $v_M \in (W^J)_{\text{af}}$.*

Proof. Let $X = g_m g_{m-1} \cdots g_2 g_1$, where $g_k \in \{e_i, f_i \mid i \in I_{\text{af}}\}$ for each $1 \leq k \leq m$. We show the assertion by induction on m . If $m = 0$, then the assertion is obvious since $X = \text{id}$. Assume that $m > 0$. Set $Y := g_{m-1} \cdots g_2 g_1$. Since $X\eta \neq \mathbf{0}$, it follows that $Y\eta \neq \mathbf{0}$. By Lemma 7.1.3, we can write $\sigma_N(Y\eta)$ as $\sigma_N(Y\eta) = [u_1 T_{\zeta_1}, u_2 T_{\zeta_2}, \dots, u_N T_{\zeta_N}]$ for some $u_1, u_2, \dots, u_N \in W_{\text{af}}$ such that $u_M T_{\zeta_M} = u_M z_{\zeta_M} t_{\zeta_M} \in (W^J)_{\text{af}}$, $1 \leq M \leq N$. By our induction hypothesis, we have $Y\eta' \neq \mathbf{0}$, and $\sigma_N(Y\eta') = [u_1 T_{\zeta'_1}, u_2 T_{\zeta'_2}, \dots, u_N T_{\zeta'_N}]$. Now, let us take $i \in I_{\text{af}}$ for which $g_m = e_i$ or f_i . We see from the definition of the root operator e_i or f_i that $\sigma_N(X\eta) = \sigma_N(g_m Y\eta)$ is of the form:

$$\sigma_N(X\eta) = [u_1 T_{\zeta_1}, \dots, u_L T_{\zeta_L}, r_i u_{L+1} T_{\zeta_{L+1}}, \dots, r_i u_K T_{\zeta_K}, u_{K+1} T_{\zeta_{K+1}}, \dots, u_N T_{\zeta_N}]$$

for some $0 \leq L < K \leq N$, with $r_i u_M T_{\zeta_M} \in (W^J)_{\text{af}}$ for all $L+1 \leq M \leq K$; remark that L and K are determined by the function $H_i^{\overline{Y\eta}}(t) = \langle \alpha_i^\vee, \overline{Y\eta}(t) \rangle$ (see the definitions of $t_0, t_1 \in [0, 1]$ in Definition 3.1.4). Because

$$\langle \alpha_i^\vee, u_M T_{\zeta'_M} \lambda \rangle = \langle \alpha_i^\vee, u_M \lambda \rangle = \langle \alpha_i^\vee, u_M T_{\zeta_M} \lambda \rangle \quad \text{for all } 1 \leq M \leq N,$$

we deduce that $H_i^{\overline{Y\eta}}(t) = H_i^{\overline{Y\eta'}}(t)$ for all $t \in [0, 1]$, which implies that t_0, t_1 for $Y\eta'$ coincide with those for $Y\eta$. Therefore, it follows from the definition of the root operator e_i or f_i that $X\eta' = g_m Y\eta' \neq \mathbf{0}$, and that

$$\begin{aligned} \sigma_N(X\eta') &= \sigma_N(g_m Y\eta') \\ &= [u_1 T_{\zeta'_1}, \dots, u_L T_{\zeta'_L}, r_i u_{L+1} T_{\zeta'_{L+1}}, \dots, r_i u_K T_{\zeta'_K}, u_{K+1} T_{\zeta'_{K+1}}, \dots, u_N T_{\zeta'_N}]. \end{aligned}$$

This proves the lemma. $\square$

Proof of Proposition 7.1.2. First, we prove that each connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ contains an element of the form (7.1.1) with $\xi_s = 0$. Let $\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$; recall that $\overline{\eta} \in \mathbb{B}(\lambda)$ by Proposition 3.1.3. From [NS4, Theorem 3.1 (2)], we know that there exists a monomial X in the root operators e_i and f_i , $i \in I_{\text{af}}$, such that $(X\overline{\eta})(t) \equiv t\lambda \pmod{\mathbb{R}\delta}$ for $t \in [0, 1]$. Because the map $\overline{} : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \rightarrow \mathbb{B}(\lambda)$ is a strict morphism of crystals (see Remark 4.3.1 (1)), it follows immediately that $X\eta \neq \mathbf{0}$, and that $X\eta \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ is of the form:

$$X\eta = (T_{\xi'_1}, T_{\xi'_2}, \dots, T_{\xi'_{s-1}}, T_\xi; a_0, a_1, \dots, a_{s-1}, a_s)$$

for some $\xi'_1, \xi'_2, \dots, \xi'_{s-1}, \xi \in Q^{\vee, J\text{-ad}}$.

Since $\eta_{T_\xi} = (T_\xi; 0, 1) \in \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ by Remark 5.1.2, there exists a monomial Y in the root operators e_i and f_i , $i \in I_{\text{af}}$, such that $Y\eta_{T_\xi} = \eta_e$. Let N be a multiple of N_λ (see the comment preceding Lemma 5.2.3). Then, $\sigma_N(\eta_{T_\xi}) = [T_\xi, \dots, T_\xi] \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ and $\sigma_N(\eta_e) = [e, \dots, e] \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$. Also, we see from Lemma 7.1.3 that $\sigma_N(Y\eta_{T_\xi}) = [v_1 T_\xi, v_2 T_\xi, \dots, v_N T_\xi]$ for some $v_1, v_2, \dots, v_N \in W_{\text{af}}$ such that $v_M T_\xi \in (W^J)_{\text{af}}$ for all $1 \leq M \leq N$. Since $Y\eta_{T_\xi} = \eta_e$, and hence

$\sigma_N(Y\eta_{T_\xi}) = \sigma_N(\eta_e)$, we deduce that $v_M T_\xi = e$ for all $1 \leq M \leq N$. Thus, we obtain $v_M = T_\xi^{-1}$ for all $1 \leq M \leq N$.

It follows that $\sigma_N(X\eta)$ is of the form $\sigma_N(X\eta) = [T_{\zeta'_1}, T_{\zeta'_2}, \dots, T_{\zeta'_N}]$ for some $\zeta'_1, \zeta'_2, \dots, \zeta'_N \in Q^{\vee, J\text{-ad}}$ with $\zeta'_N = \xi$. Therefore, from Lemma 7.1.4 applied to η_{T_ξ} and $X\eta$, we deduce that

$$\sigma_N(YX\eta) = [v_1 T_{\zeta'_1}, v_2 T_{\zeta'_2}, \dots, v_N T_{\zeta'_N}] = [T_\xi^{-1} T_{\zeta'_1}, T_\xi^{-1} T_{\zeta'_2}, \dots, \underbrace{T_\xi^{-1} T_{\zeta'_N}}_{=e}],$$

where $v_M T_{\zeta'_M} = T_\xi^{-1} T_{\zeta'_M} \in (W^J)_{\text{af}}$ for all $1 \leq M \leq N$; using (2.3.5), we see by direct computation that $T_\xi^{-1} T_{\zeta'_M} \in (W^J)_{\text{af}}$ is of the form:

$$T_\xi^{-1} T_{\zeta'_M} = z_{\zeta_M} t_{\zeta_M} = T_{\zeta_M} \quad \text{for some } \zeta_M \in Q^{\vee, J\text{-ad}}. \quad (7.1.2)$$

Hence we obtain $\sigma_N(YX\eta) = [T_{\zeta_1}, T_{\zeta_2}, \dots, T_{\zeta_{N-1}}, e]$. Because the final factor of $\sigma_N(YX\eta)$ is identical to $e = T_0$, we conclude that $YX\eta$ is of the form (7.1.1) with $\xi_s = 0$. Thus, we have proved that each connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ contains an element of the form (7.1.1) with $\xi_s = 0$.

Next, we prove the uniqueness statement. Let $\eta, \eta' \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, $\eta \neq \eta'$, be of the form (7.1.1) with $\xi_s = 0$, and suppose that $X\eta = \eta'$ for some monomial X in the root operators e_i and f_i , $i \in I_{\text{af}}$. As above, let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ . Then, $\sigma_N(\eta)$ and $\sigma_N(\eta')$ are of the form (note that $T_0 = e$):

$$\sigma_N(\eta) = [T_{\zeta_1}, T_{\zeta_2}, \dots, T_{\zeta_{N-1}}, T_0], \quad \sigma_N(\eta') = [T_{\zeta'_1}, T_{\zeta'_2}, \dots, T_{\zeta'_{N-1}}, T_0],$$

for some $\zeta_M, \zeta'_M \in Q^{\vee, J\text{-ad}}$, $1 \leq M \leq N-1$, respectively. Since $\eta \neq \eta'$ and the map $\sigma_N : \mathbb{B}^{\frac{\infty}{2}}(\lambda) \hookrightarrow \mathbb{B}^{\frac{\infty}{2}}(\lambda)^{\otimes N}$ is injective, there exists $1 \leq M \leq N-1$ such that $\zeta_M \neq \zeta'_M$; let $1 \leq L \leq N-1$ be the maximum of all such M 's. Then we deduce from Lemma 2.3.5(1) that $\zeta'_L - \zeta_L \notin Q_J^\vee$, and hence $[\zeta'_L - \zeta_L] \neq 0$; here, by interchanging η and η' if necessary, we may assume that $[\zeta'_L - \zeta_L] \notin Q_{J^c}^{\vee+}$.

From Lemma 7.1.3, we have

$$\sigma_N(X\eta) = [v_1 T_{\zeta_1}, v_2 T_{\zeta_2}, \dots, v_{N-1} T_{\zeta_{N-1}}, v_N T_0]$$

for some $v_1, v_2, \dots, v_N \in W_{\text{af}}$ such that $v_M T_{\zeta_M} \in (W^J)_{\text{af}}$, $1 \leq M \leq N-1$, and $v_N T_0 \in (W^J)_{\text{af}}$; note that $v_1, v_2, \dots, v_N \in (W^J)_{\text{af}}$ by Lemma 2.3.7. Since $X\eta = \eta'$ by our assumption, it follows that $v_M T_{\zeta_M} = T_{\zeta'_M}$ for all $1 \leq M \leq N-1$, and $v_N T_0 = T_0$. Therefore, we obtain $v_M = T_{\zeta'_M} T_{\zeta_M}^{-1}$ for all $1 \leq M \leq N-1$, and $v_N = e$; in particular, $v_{L+1} = \dots = v_{N-1} = v_N = e$ by the definition of L . Also, the same reasoning as for (7.1.2) shows that for each $1 \leq M \leq N$, $v_M = T_{\zeta'_M} T_{\zeta_M}^{-1} \in (W^J)_{\text{af}}$ is of the form $v_M = T_{\zeta''_M}$ for some $\zeta''_M \in Q^{\vee, J\text{-ad}}$ such that $[\zeta''_M] = [\zeta'_M - \zeta_M]$. Therefore, from Lemma 7.1.4 applied to η and η_e , we deduce that $X\eta_e \neq \mathbf{0}$, and that

$$\sigma_N(X\eta_e) = [v_1, v_2, \dots, v_N] = [T_{\zeta''_1}, T_{\zeta''_2}, \dots, T_{\zeta''_L}, e, \dots, e],$$

which implies that $X\eta_e \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ is of the form:

$$X\eta_e = (T_{\xi_1}, \dots, T_{\xi_{s-1}}, e; a_0, a_1, \dots, a_{s-1}, a_s),$$

for some $\xi_1, \dots, \xi_{s-1} \in Q^{\vee, J\text{-ad}}$, with $\xi_{s-1} = \zeta_L''$. Hence we obtain $[\xi_{s-1} - 0] = [\zeta_L''] = [\zeta_L' - \zeta_L] \notin Q_{J^c}^{\vee+}$, which contradicts Proposition 7.1.1. This completes the proof of Proposition 7.1.2. $\square$

7.2 Proof of Proposition 3.2.4.

Recall that $\text{Turn}(\lambda) = \{k/m_i \mid i \in J^c = I \setminus J, 0 \leq k \leq m_i\}$; we enumerate the elements of $\text{Turn}(\lambda)$ in increasing order as:

$$\text{Turn}(\lambda) = \{0 = \tau_0 < \tau_1 < \dots < \tau_p = 1\}.$$

Let $0 \leq q \leq p$. Note that $i \in J^c(\lambda; \tau_q)$ if and only if $i \notin J$ and $\tau_q = k/m_i$ for some $0 \leq k \leq m_i$.

Proposition 7.2.1. *There exists a bijective correspondence between the set $\text{Par}(\lambda)$ and the set of all connected components of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$.*

Proof. Let $\text{Conn}(\lambda)$ denote the set of all connected components of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$. First, we define a map $\Theta : \text{Conn}(\lambda) \rightarrow \text{Par}(\lambda)$ as follows. Take an arbitrary $C \in \text{Conn}(\lambda)$. By Propositions 7.1.1 and 7.1.2, the connected component C contains a unique element η of the form $\eta = (T_{\xi_1}, T_{\xi_2}, \dots, T_{\xi_{s-1}}, e; a_0, a_1, \dots, a_{s-1}, a_s)$, with $\xi_1, \xi_2, \dots, \xi_{s-1} \in Q^{\vee, J\text{-ad}}$, such that $a_u \in \text{Turn}(\lambda)$ for all $0 \leq u \leq s$, and such that $[\xi_u - \xi_{u+1}] \in Q_{J^c(\lambda; a_u)}^{\vee+}$ for all $1 \leq u \leq s-1$, where we set $\xi_s := 0$. For each $0 \leq u \leq s$, let $0 \leq q_u \leq p$ be such that $a_u = \tau_{q_u}$. Then we define ζ_q , $1 \leq q \leq p$, by $\zeta_q := \xi_u$ if $q_{u-1} + 1 \leq q \leq q_u$, that is,

$$\underbrace{\zeta_1, \dots, \zeta_{q_1}}_{:=\xi_1}, \underbrace{\zeta_{q_1+1}, \dots, \zeta_{q_2}}_{:=\xi_2}, \dots, \underbrace{\zeta_{q_{s-2}+1}, \dots, \zeta_{q_{s-1}}}_{:=\xi_{s-1}}, \underbrace{\zeta_{q_{s-1}+1}, \dots, \zeta_p}_{:=\xi_s=0};$$

remark that for all $1 \leq q \leq p-1$,

$$[\zeta_q - \zeta_{q+1}] \in Q_{J^c(\lambda; \tau_q)}^{\vee+}. \quad (7.2.1)$$

Fix $i \in J^c = I \setminus J$, and let $c_q^{(i)}$, $1 \leq q \leq p$, be the coefficient of $\alpha_i^{\vee}$ in ζ_q ; we see from (7.2.1) that $c_1^{(i)} \geq c_2^{(i)} \geq \dots \geq c_{p-1}^{(i)} \geq c_p^{(i)} = 0$, and that $c_q^{(i)} = c_{q+1}^{(i)}$ for $1 \leq q \leq p-1$ such that $i \notin J^c(\lambda; \tau_q)$, i.e., $\tau_q \notin \{k/m_i \mid 0 \leq k \leq m_i\}$. For each $1 \leq k \leq m_i$, let $1 \leq p_k \leq p$ be such that $\tau_{p_k} = k/m_i$. Then we define $\rho_k^{(i)}$, $1 \leq k \leq m_i - 1$, in the following way:

$$\begin{aligned} \underbrace{c_1^{(i)} = \dots = c_{p_1}^{(i)}}_{=:\rho_1^{(i)}} &\geq \underbrace{c_{p_1+1}^{(i)} = \dots = c_{p_2}^{(i)}}_{=:\rho_2^{(i)}} \geq \dots \\ &\dots \geq \underbrace{c_{p_{m_i-2}+1}^{(i)} = \dots = c_{p_{m_i-1}}^{(i)}}_{=:\rho_{m_i-1}^{(i)}} \geq c_{p_{m_i-1}+1}^{(i)} = \dots = c_{p_{m_i}}^{(i)} = 0. \end{aligned}$$

Thus we obtain a partition $\rho^{(i)} := (\rho_1^{(i)} \geq \rho_2^{(i)} \geq \cdots \geq \rho_{m_i-1}^{(i)})$ of length less than m_i . Now we define $\Theta(C) := (\rho^{(i)})_{i \in I} \in \text{Par}(\lambda)$, where for every $j \in J$, we define $\rho^{(j)}$ to be the empty partition.

Next, we define a map $\Xi : \text{Par}(\lambda) \rightarrow \text{Conn}(\lambda)$ as follows. Take an arbitrary $\boldsymbol{\rho} = (\rho^{(i)}) \in \text{Par}(\lambda)$, with $\rho^{(i)} = (\rho_1^{(i)} \geq \rho_2^{(i)} \geq \cdots \geq \rho_{m_i-1}^{(i)})$ for $i \in I \setminus J$. Define $\zeta_q \in Q^\vee$, $1 \leq q \leq p$, inductively by

$$\zeta_p := 0, \quad \text{and} \quad \zeta_q := \zeta_{q+1} + \sum_{i \in J^c(\lambda; \tau_q)} (\rho_{\tau_q m_i}^{(i)} - \rho_{\tau_q m_i + 1}^{(i)}) \alpha_i^\vee \quad \text{for } 1 \leq q \leq p-1;$$

note that for $1 \leq q \leq p-1$, if $i \in J^c(\lambda; \tau_q)$, then $\tau_q m_i = \tau_q \langle \alpha_i^\vee, \lambda \rangle \in \mathbb{Z}$, with $1 \leq \tau_q m_i \leq m_i - 1$. We write the set $\{1 \leq q \leq p-1 \mid \zeta_q \neq \zeta_{q+1}\}$ in the form $\{q_1 < q_2 < \cdots < q_{s-1}\}$, i.e.,

$$\zeta_1 = \cdots = \zeta_{q_1} \neq \zeta_{q_1+1} = \cdots = \zeta_{q_2} \neq \cdots \neq \zeta_{q_{s-1}+1} = \cdots = \zeta_p = 0.$$

Then we define (for the definition of $\phi_J(\zeta_{p_u})$, see Lemma 2.3.5)

$$\begin{cases} \xi_s := 0, & \xi_u := \zeta_{q_u} + \phi_J(\zeta_{q_u}) \quad \text{for } 1 \leq u \leq s-1, \\ a_0 := 0, & a_u := \tau_{q_u} \quad \text{for } 1 \leq u \leq s-1, \quad a_s := 1; \end{cases}$$

it follows from Proposition 7.1.1 that

$$\eta_{\boldsymbol{\rho}} := (T_{\xi_1}, T_{\xi_2}, \dots, T_{\xi_{s-1}}, e; a_0, a_1, \dots, a_{s-1}, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda). \quad (7.2.2)$$

Now we define $\Xi(\boldsymbol{\rho})$ to be the connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ containing this $\eta_{\boldsymbol{\rho}}$.

We deduce from the definitions that the maps Θ and Ξ are inverses of each other. This completes the proof of the proposition. $\square$

Proof of Proposition 3.2.4. For $\boldsymbol{\rho} \in \text{Par}(\lambda)$, let $\eta_{\boldsymbol{\rho}} \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ be as defined by (7.2.2), which is a unique element of the form (7.1.1) with $\xi_s = 0$ contained in the connected component $\mathbb{B}_{\boldsymbol{\rho}}^{\frac{\infty}{2}}(\lambda) := \Xi(\boldsymbol{\rho})$. We prove that there exists a unique isomorphism $\mathbb{B}_{\boldsymbol{\rho}}^{\frac{\infty}{2}}(\lambda) \xrightarrow{\sim} \{\boldsymbol{\rho}\} \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals that maps $\eta_{\boldsymbol{\rho}}$ to $\boldsymbol{\rho} \otimes \eta_e$. As in the proof of Proposition 3.2.2, it suffices to show the following for monomials X, Y in the Kashiwara operators:

- (i) $X\eta_{\boldsymbol{\rho}} \neq \mathbf{0}$ if and only if $X(\boldsymbol{\rho} \otimes \eta_e) \neq \mathbf{0}$;
- (ii) $X\eta_{\boldsymbol{\rho}} = Y\eta_{\boldsymbol{\rho}}$ if and only if $X(\boldsymbol{\rho} \otimes \eta_e) = Y(\boldsymbol{\rho} \otimes \eta_e)$.

Part (i) follows immediately from Lemma 7.1.4 and the equality $X(\boldsymbol{\rho} \otimes \eta_e) = \boldsymbol{\rho} \otimes X\eta_e$. Let us show part (ii). We give a proof only for the “only if” part; the proof for the “if” part is similar. Assume that $X\eta_{\boldsymbol{\rho}} = Y\eta_{\boldsymbol{\rho}} \neq \mathbf{0}$. Let $N \in \mathbb{Z}_{>0}$ be a multiple of N_λ (see the comment preceding Lemma 5.2.3), and write $\sigma_N(\eta_{\boldsymbol{\rho}})$ as $\sigma_N(\eta_{\boldsymbol{\rho}}) = [T_{\xi_1}, T_{\xi_2}, \dots, T_{\xi_N}]$ for some $\xi_1, \xi_2, \dots, \xi_N \in Q^\vee, J\text{-ad}$. By Lemma 7.1.3, $\sigma_N(X\eta_{\boldsymbol{\rho}})$ and $\sigma_N(Y\eta_{\boldsymbol{\rho}})$ are of the form:

$$\sigma_N(X\eta_{\boldsymbol{\rho}}) = [u_1 T_{\xi_1}, u_2 T_{\xi_2}, \dots, u_N T_{\xi_N}], \quad \sigma_N(Y\eta_{\boldsymbol{\rho}}) = [v_1 T_{\xi_1}, v_2 T_{\xi_2}, \dots, v_N T_{\xi_N}],$$

for some $u_1, u_2, \dots, u_N \in (W^J)_{\text{af}}$ and $v_1, v_2, \dots, v_N \in (W^J)_{\text{af}}$, respectively. Then, by Lemma 7.1.4, we have

$$\sigma_N(X\eta_e) = [u_1, u_2, \dots, u_N], \quad \sigma_N(Y\eta_e) = [v_1, v_2, \dots, v_N],$$

respectively. Since $X\eta_\rho = Y\eta_\rho$ by the assumption, we have $u_M = v_M$ for all $1 \leq M \leq N$. Therefore, we see that $\sigma_N(X\eta_e) = \sigma_N(Y\eta_e)$, and hence $X\eta_e = Y\eta_e$ by the injectivity of σ_N . Thus, we obtain $X(\rho \otimes \eta_e) = \rho \otimes X\eta_e = \rho \otimes Y\eta_e = Y(\rho \otimes \eta_e)$, as desired. Finally, from the existence of the isomorphism $\mathbb{B}_\rho^{\frac{\infty}{2}}(\lambda) \xrightarrow{\sim} \{\rho\} \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals for each $\rho \in \text{Par}(\lambda)$, together with Proposition 7.2.1, we conclude that

$$\mathbb{B}^{\frac{\infty}{2}}(\lambda) = \bigsqcup_{\rho \in \text{Par}(\lambda)} \mathbb{B}_\rho^{\frac{\infty}{2}}(\lambda) \cong \bigsqcup_{\rho \in \text{Par}(\lambda)} \{\rho\} \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda) = \text{Par}(\lambda) \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda).$$

This completes the proof of Proposition 3.2.4. $\square$

Remark 7.2.2. Since $\mathcal{B}(\lambda) \cong \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ as crystals, we can define the action of the Weyl group W_{af} on $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ by the same formula as (5.1.1) for the one on $\mathcal{B}(\lambda)$. Let $\rho \in \text{Par}(\lambda)$. Then, as shown in the proof of Proposition 3.2.4, there exists a unique isomorphism $\mathbb{B}_\rho^{\frac{\infty}{2}}(\lambda) \xrightarrow{\sim} \{\rho\} \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals that maps η_ρ to $\rho \otimes \eta_e$. Hence it follows from the tensor product rule for crystals that $x(\rho \otimes \eta_e) = \rho \otimes (x\eta_e)$ for every $x \in W_{\text{af}}$. From this equality, again using the tensor product rule for crystals, we deduce that η_ρ is an extremal element of weight $\lambda + \text{wt}(\rho)$, since η_e is an extremal element of weight λ by Remark 5.3.2.

A Appendix.

A.1 Relation between the semi-infinite Bruhat graph and the quantum Bruhat graph.

In this subsection, we fix a subset $J \subset I$. Set $\rho_J := \frac{1}{2} \sum_{\alpha \in \Delta_J^+} \alpha$; note that

$$\langle \xi, \rho - \rho_J \rangle = 0 \quad \text{for all } \xi \in Q_J^\vee. \quad (\text{A.1.1})$$

Definition A.1.1 ([LNS³1, §4]; see also [BFP, §6]). Let J be a subset of I . Define the (parabolic) quantum Bruhat graph QB^J to be the $(\Delta^+ \setminus \Delta_J^+)$ -labeled, directed graph with vertex set W^J and $(\Delta^+ \setminus \Delta_J^+)$ -labeled, directed edges of the following form: $w \xrightarrow{\gamma} \lfloor wr_\gamma \rfloor$ for $w \in W^J$ and $\gamma \in \Delta^+ \setminus \Delta_J^+$ such that either (i) $\ell(\lfloor wr_\gamma \rfloor) = \ell(w) + 1$, or (ii) $\ell(\lfloor wr_\gamma \rfloor) = \ell(w) + 1 - 2\langle \gamma^\vee, \rho - \rho_J \rangle$; we call an edge $w \xrightarrow{\gamma} \lfloor wr_\gamma \rfloor$ satisfying condition (i) (resp., (ii)) a Bruhat (resp., quantum) edge.

Combining Proposition 4.2.1 and [LNS³1, Theorem 6.5], we obtain the following.

Proposition A.1.2.

- (1) Let $x = wz_\xi t_\xi \in (W^J)_{\text{af}}$ with $w \in W^J$ and $\xi \in Q^{\vee, J\text{-ad}}$, and $\beta \in \Delta_{\text{af}}^+$. Assume that $x \xrightarrow{\beta} r_\beta x$ in SiB^J ; note that $\beta = w\gamma + n\delta$ for some $\gamma \in \Delta^+ \setminus \Delta_J^+$ and $n \in \{0, 1\}$ by Corollary 4.2.2. If $n = 0$ (resp., $n = 1$), then we have a Bruhat edge (reps., quantum edge) $w \xrightarrow{\gamma} \lfloor wr_\gamma \rfloor$ in QB^J .
- (2) Let $w \in W^J$, and $\gamma \in \Delta^+ \setminus \Delta_J^+$. Assume that $w \xrightarrow{\gamma} \lfloor wr_\gamma \rfloor$ in QB^J . Set $\beta := w\gamma$ (resp., $\beta := w\gamma + \delta$) if the edge is a Bruhat edge (resp., quantum edge). Then, $\beta \in \Delta_{\text{af}}^+$, and $wz_\xi t_\xi \xrightarrow{\beta} r_\beta wz_\xi t_\xi$ in SiB^J for every $\xi \in Q^{\vee, J\text{-ad}}$.

A.2 Another definition of the semi-infinite Bruhat order.

In this subsection, we fix a subset $J \subset I$. For $vt_\zeta \in W_{\text{af}}$ with $v \in W$ and $\zeta \in Q^\vee$, we define

$$\ell_J^{\frac{\infty}{2}}(vt_\zeta) := \ell(\lfloor v \rfloor) + 2\langle \zeta, \rho - \rho_J \rangle. \quad (\text{A.2.1})$$

Lemma A.2.1. *The equalities $\ell_J^{\frac{\infty}{2}}(x) = \ell_J^{\frac{\infty}{2}}(\Pi^J(x)) = \ell^{\frac{\infty}{2}}(\Pi^J(x))$ hold for all $x \in W_{\text{af}}$.*

Proof. We write $\Pi^J(x)$ as $\Pi^J(x) = wz_\xi t_\xi$, with $w \in W^J$ and $\xi \in Q^{\vee, J\text{-ad}}$. The second equality follows from [LNS³1, (3.11)] and the equality $\ell(wz_\xi) = \ell(w) + \ell(z_\xi)$.

In order to prove the first equality, we write $x = x_1 x_2$ with $x_1 \in (W^J)_{\text{af}}$ and $x_2 \in (W_J)_{\text{af}}$; note that $\Pi^J(x) = x_1$. We have $x_1 = w_1 z_{\xi_1} t_{\xi_1}$ for some $w_1 \in W^J$ and $\xi_1 \in Q^{\vee, J\text{-ad}}$ by (2.3.5), and $x_2 = w_2 t_{\xi_2}$ for some $w_2 \in W_J$ and $\xi_2 \in Q_J^\vee$ by (2.3.3). Since $x = x_1 x_2 = w_1 z_{\xi_1} w_2 t_{w_2^{-1}\xi_1 + \xi_2}$, we compute

$$\begin{aligned} \ell_J^{\frac{\infty}{2}}(x) &= \ell(\lfloor w_1 z_{\xi_1} w_2 \rfloor) + 2\langle w_2^{-1}\xi_1 + \xi_2, \rho - \rho_J \rangle \\ &= \ell(w_1) + 2\langle w_2^{-1}\xi_1 + \xi_2, \rho - \rho_J \rangle \quad \text{since } w_1 \in W^J \text{ and } z_{\xi_1} w_2 \in W_J \\ &= \ell(w_1) + 2\langle \xi_1, \rho - \rho_J \rangle \quad \text{since } \xi_2 \in Q_J^\vee \text{ and } w_2 \in W_J \text{ (see (A.1.1))} \\ &= \ell_J^{\frac{\infty}{2}}(w_1 z_{\xi_1} t_{\xi_1}) = \ell_J^{\frac{\infty}{2}}(x_1) = \ell_J^{\frac{\infty}{2}}(\Pi^J(x)). \end{aligned}$$

This proves the lemma. □

Proposition A.2.2. *Let $x, y \in (W^J)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$. We have $x \xrightarrow{\beta} y$ in SiB^J if and only if the following three conditions are satisfied:*

- (a) $y = \Pi^J(r_\beta x)$;
- (b) $\ell_J^{\frac{\infty}{2}}(r_\beta x) = \ell_J^{\frac{\infty}{2}}(x) + 1$;
- (c) if we write x as $x = wz_\xi t_\xi$ with $w \in W^J$ and $\xi \in Q^{\vee, J\text{-ad}}$, then $\beta = w\gamma + n\delta$ for some $\gamma \in \Delta^+ \setminus \Delta_J^+$ and $n \in \{0, 1\}$.

Proof. The “only if” part follows immediately from Corollary 4.2.2 and Lemma A.2.1. We show the “if” part. By condition (c), we have $r_\beta x = wr_\gamma z_\xi t_{\xi+nz_\xi^{-1}\gamma^\vee}$. We compute

$$\begin{aligned} 1 &= \ell_J^{\frac{\infty}{2}}(r_\beta x) - \ell_J^{\frac{\infty}{2}}(x) \quad \text{by (b)} \\ &= \ell(\lfloor wr_\gamma \rfloor) + 2\langle \xi + nz_\xi^{-1}\gamma^\vee, \rho - \rho_J \rangle - \ell(w) - 2\langle \xi, \rho - \rho_J \rangle \\ &= \ell(\lfloor wr_\gamma \rfloor) - \ell(w) + 2n\langle \gamma^\vee, \rho - \rho_J \rangle \quad (\text{see (A.1.1)}). \end{aligned}$$

From this, using the condition that $n \in \{0, 1\}$, we deduce that $w \xrightarrow{\gamma} \lfloor wr_\gamma \rfloor$ in QB^J ; observe that this edge is a Bruhat (resp., quantum) edge if and only if $n = 0$ (resp., $n = 1$). Therefore, by Proposition A.1.2 (2), we have $x = wz_\xi t_\xi \xrightarrow{\beta} r_\beta wz_\xi t_\xi = r_\beta x$ in SiB^J ; in particular, $r_\beta x \in (W^J)_{\text{af}}$. Thus, by condition (a), we obtain $y = \Pi^J(r_\beta x) = r_\beta x$, and hence $x \xrightarrow{\beta} y$ in SiB^J . This proves the proposition. $\square$

A.3 Relation between the semi-infinite Bruhat order and the generic Bruhat order.

In this subsection, we assume that $J = \emptyset$; note that $(W^J)_{\text{af}} = W_{\text{af}}$. Fix an (arbitrary) element $\xi \in Q^\vee$ such that $\langle \xi, \alpha_i \rangle > 0$ for all $i \in I$. We know from [P] (see also [LNS³1, Theorem 5.2], and Proposition A.1.2) that for $x, y \in (W^J)_{\text{af}} = W_{\text{af}}$, $x \leq_{\frac{\infty}{2}} y$ if and only if there exists $N \in \mathbb{Z}_{\geq 0}$, depending on x, y , and ξ , such that $yt_{-n\xi} \leq xt_{-n\xi}$ (or equivalently, $t_{n\xi}y^{-1} \leq t_{n\xi}x^{-1}$) for all $n \in \mathbb{Z}_{\geq N}$, where $\leq$ is the (ordinary) Bruhat order on W_{af} . Also, in [Lu1, §1.5], Lusztig introduced a partial order $\leq_L$ on W_{af} , which we call Lusztig’s generic Bruhat order; we know from [S, Claim 4.14 in the proof of Lemma 4.13] that $x \leq_L y$ if and only if there exists $N \in \mathbb{Z}_{\geq 0}$, depending on x, y , and ξ , such that $t_{n\xi}x \leq t_{n\xi}y$ for all $n \in \mathbb{Z}_{\geq N}$. Combining these facts, we obtain the following.

Lemma A.3.1. *Let $x, y \in W_{\text{af}}$. We have $x \leq_{\frac{\infty}{2}} y$ if and only if $y^{-1} \leq_L x^{-1}$.*

References

- [AK] T. Akasaka and M. Kashiwara, Finite-dimensional representations of quantum affine algebras, *Publ. Res. Inst. Math. Sci.* **33** (1997), 839–867.
- [BN] J. Beck and H. Nakajima, Crystal bases and two-sided cells of quantum affine algebras, *Duke Math. J.* **123** (2004), 335–402.
- [BB] A. Björner and F. Brenti, “Combinatorics of Coxeter Groups”, Graduate Texts in Mathematics Vol. 231, Springer, New York, 2005.
- [BF] A. Braverman and M. Finkelberg, Weyl modules and q -Whittaker functions, *Math. Ann.* **359** (2014), 45–59.
- [BFP] F. Brenti, S. Fomin, and A. Postnikov, Mixed Bruhat operators and Yang-Baxter equations for Weyl groups, *Int. Math. Res. Not.* **8** (1999), 419–441.

- [CP] V. Chari and A. Pressley, Weyl modules for classical and quantum affine algebras, *Represent. Theory* **5** (2001), 191–223.
- [FFKM] B. Feigin, M. Finkelberg, A. Kuznetsov, and I. Mirković, Semi-infinite flags II: Local and global intersection cohomology of quasimaps’ spaces, *Differential Topology, Infinite-Dimensional Lie Algebras, and Applications*, Amer. Math. Soc. Transl. Ser. 2, Vol. 194, pp. 113–148, Amer. Math. Soc., Providence, RI, 1990.
- [FF] B. Feigin and E. Frenkel, Affine Kac-Moody algebras and semi-infinite flag manifolds, *Comm. Math. Phys.* **128** (1990), 161–189.
- [HN] D. Hernandez and H. Nakajima, Level 0 monomial crystals, *Nagoya Math. J.* **184** (2006), 85–153.
- [HK] J. Hong and S.-J. Kang, “Introduction to Quantum Groups and Crystal Bases”, Graduate Studies in Mathematics Vol. 42, Amer. Math. Soc., Providence, RI, 2002.
- [Kac] V. G. Kac, “Infinite Dimensional Lie Algebras”, 3rd Edition, Cambridge University Press, Cambridge, UK, 1990.
- [Kas1] M. Kashiwara, Crystal bases of modified quantized enveloping algebra, *Duke Math. J.* **73** (1994), 383–413.
- [Kas2] M. Kashiwara, On crystal bases, in “Representations of Groups” (B.N. Allison and G.H. Cliff, Eds.), CMS Conf. Proc. Vol. 16, pp. 155–197, Amer. Math. Soc., Providence, RI, 1995.
- [Kas3] M. Kashiwara, Similarity of crystal bases, in “Lie Algebras and Their Representations” (S.-J. Kang et al., Eds.), Contemp. Math. Vol. 194, pp. 177–186, Amer. Math. Soc., Providence, RI, 1996.
- [Kas4] M. Kashiwara, “Bases Cristallines des Groupes Quantiques” (Notes by Charles Cochet), Cours Spécialisés Vol. 9, Société Mathématique de France, Paris, 2002.
- [Kas5] M. Kashiwara, On level-zero representations of quantized affine algebras, *Duke Math. J.* **112** (2002), 117–175.
- [LS] T. Lam and M. Shimozono, Quantum cohomology of G/P and homology of affine Grassmannian, *Acta Math.* **204** (2010), 49–90.
- [LNS³1] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals I: lifting the parabolic quantum Bruhat graph, arXiv:1211.2042v3, to appear in *Int. Math. Res. Not.*
- [LNS³2] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals II: Alcove model, path model, and $P = X$, preprint 2014, arXiv:1402.2203.
- [Li1] P. Littelmann, A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras, *Invent. Math.* **116** (1994), 329–346.
- [Li2] P. Littelmann, Paths and root operators in representation theory, *Ann. of Math.* (2) **142** (1995), 499–525.
- [Lu1] G. Lusztig, Hecke algebras and Jantzen’s generic decomposition patterns, *Adv. Math.* **37** (1980), 121–164.

- [Lu2] G. Lusztig, Canonical bases in tensor products, *Proc. Nat. Acad. Sci. U.S.A.* **89** (1992), 8177–8179.
- [Mi] L. C. Mihalcea, On equivariant quantum cohomology of homogeneous spaces: Chevalley formulae and algorithms, *Duke Math. J.* **140** (2007), 321–350.
- [NS1] S. Naito and D. Sagaki, Path model for a level-zero extremal weight module over a quantum affine algebra, *Int. Math. Res. Not.* **2003** (2003), no. 32, 1731–1754.
- [NS2] S. Naito and D. Sagaki, Crystal of Lakshmibai-Seshadri paths associated to an integral weight of level zero for an affine Lie algebra, *Int. Math. Res. Not.* **2005** (2005), no. 14, 815–840.
- [NS3] S. Naito and D. Sagaki, Path model for a level-zero extremal weight module over a quantum affine algebra. II, *Adv. Math.* **200** (2006), 102–124.
- [NS4] S. Naito and D. Sagaki, Crystal structure on the set of Lakshmibai-Seshadri paths of an arbitrary level-zero shape, *Proc. Lond. Math. Soc.* (3) **96** (2008), 582–622.
- [NS5] S. Naito and D. Sagaki, Demazure submodules of level-zero extremal weight modules and specializations of Macdonald polynomials, preprint 2014, arXiv:1404.2436v2.
- [N1] H. Nakajima, Quiver varieties and finite-dimensional representations of quantum affine algebras, *J. Amer. Math. Soc.* **14** (2001), 145–238.
- [N2] H. Nakajima, Extremal weight modules of quantum affine algebras, in “Representation Theory of Algebraic Groups and Quantum Groups” (T. Shoji et al., Eds.), Adv. Stud. Pure Math. Vol. 40, pp. 343–369, Math. Soc. Japan, 2004.
- [P] D. Peterson, Quantum cohomology of G/P , Lecture Notes, Cambridge, MA, Spring: Massachusetts Institute of Technology, 1997.
- [S] W. Soergel, Kazhdan-Lusztig polynomials and a combinatoric for tilting modules, *Represent. Theory* **1** (1997), 83–114.